

Rabbit TYMP Polyclonal Antibody | anti-TYMP antibody

TYMP Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA36982	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TYMP; Polyclonal Antibody; TYMP Polyclonal Antibody; ECGF1; MNGIE; PDECGF; TP; hPD-ECGF; anti-TYMP antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	482
Applicable Applications for anti-TYMP antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human TYMP
Calculated Molecular Weight	50kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES


IF (Immunofluorescence)

Immunofluorescence analysis of HeLa cells using TYMP antibody. Blue: DAPI for nuclear staining.


WB (Western Blot)

Western blot analysis of extracts of various cell lines, using TYMP antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.

ADDITIONAL INFORMATION

Related Product Information for anti-TYMP antibody	Thymidine phosphorylase (TP) is a platelet-derived endothelial cell growth factor (PD-ECGF) that catalyzes the formation of thymine and 2-deoxy-D-ribose-1-phosphate from thymidine and orthophosphate (1). This intracellular enzyme is capable of both promoting angiogenesis and inhibiting apoptosis. Thymidine phosphorylase catalytic activity is required for its angiogenic function (2,3). Increased expression of TP/PD-ECGF is seen in a wide variety of different solid tumors and inflammatory diseases and is often associated with poor prognosis (4,5). Alternatively, TP can activate fluorouracil derivative (DFUR) prodrugs and increase the antitumor activity of the related treatment (1,5). The use of thymidine phosphorylase as a cancer therapeutic target has been studied extensively, with emphasis on either inhibiting TP enzymatic activity or increasing enzyme induction with concomitant DFUR treatment (1,5).
Product Categories/Family for anti-TYMP antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	67477361
NCBI GeneID	1890
UniProt Accession #	P19971; Q13390; Q8WVB7; A8MW15; H9KVA0
Molecular Weight	482
NCBI Official Full Name	Thymidine phosphorylase
NCBI Official Synonym Full Names	thymidine phosphorylase
NCBI Official Symbol	TYMP
NCBI Official Synonym Symbols	TP; ECGF; ECGF1; MNGIE; MEDPS1; MTDPS1; PDECGF; hPD-ECGF
NCBI Protein Information	thymidine phosphorylase; tdRPase; gliostatin
UniProt Protein Name	Thymidine phosphorylase
UniProt Gene Name	TYMP
UniProt Synonym Gene Names	ECGF1; TP; PD-ECGF
UniProt Entry Name	TYPH_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.