

Rabbit TP63 Polyclonal Antibody | anti-TP63 antibody

TP63 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA37105	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TP63; Polyclonal Antibody; TP63 Polyclonal Antibody; AIS; B(p51A); B(p51B); EEC3; KET; LMS; NBP; OFC8; RHS; SHFM4; TP53CP; TP53L; TP73L; p40; p51; p53CP; p63; p73H; p73L; anti-TP63 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.09% Sodium azide, 50% glycerol, pH7.3.
Concentration	1mg/ml (varies by lot)
Sequence Length	680
Applicable Applications for anti-TP63 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human TP63
Calculated Molecular Weight	77kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IF (Immunofluorescence)

Immunofluorescence analysis of U2OS cells using TP63 antibody. Blue: DAPI for nuclear staining.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human skin using TP63 antibody at dilution of 1:50 (40x lens).

ADDITIONAL INFORMATION

Related Product Information for anti-TP63 antibody	The p53 tumor suppressor protein plays a major role in cellular response to DNA damage and other genomic aberrations. Activation of p53 can lead to either cell cycle arrest and DNA repair or apoptosis (1). In addition to p53, mammalian cells contain two p53 family members, p63 and p73, which are similar to p53 in both structure and function (2). While p63 can induce p53-responsive genes and apoptosis, mutation of p63 rarely results in tumors (2). Amplification of the p63 gene is frequently observed in squamous cell carcinomas of the lung, head and neck (2,3). The p63 gene contains an alternative transcription initiation site that yields a 40 kDa deltaNp63 lacking the transactivation domain, and alternative splicing at the carboxy-terminus yields the alpha, beta and gamma isoforms (3,4).
Product Categories/Family for anti-TP63 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	57013009
NCBI GeneID	8626
UniProt Accession #	Q9H3D4; O75080; O75195; O75922; O76078; Q6VEG2; Q6VEG3; Q6VEG4; Q6VFJ1; Q6VFJ2; Q6VFJ3; Q6VH20
Molecular Weight	680
NCBI Official Full Name	Tumor protein 63
NCBI Official Synonym Full Names	tumor protein p63
NCBI Official Symbol	TP63
NCBI Official Synonym Symbols	AIS; KET; LMS; NBP; RHS; p40; p51; p63; EEC3; OFC8; p73H; p73L; SHFM4; TP53L; TP73L; p53CP; TP53CP; B(p51A); B(p51B)
NCBI Protein Information	tumor protein 63; CUSP; transformation-related protein 63; tumor protein p53-competing protein; amplified in squamous cell carcinoma; chronic ulcerative stomatitis protein; keratinocyte transcription factor KET; tumor protein p63 deltaN isoform delta
UniProt Protein Name	Tumor protein 63
UniProt Gene Name	TP63
UniProt Synonym Gene Names	KET; P63; P73H; P73L; TP73L; p63; CUSP; TP63; p73L
UniProt Entry Name	P63_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tp63-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/37105>