

Rabbit anti-Human CD86 Polyclonal Antibody | anti-CD86 antibody

CD86

CATALOG # AAA37119	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

CD86; Polyclonal Antibody; anti-CD86 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Sequence Length	217
Applicable Applications for anti-CD86 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 30-247 of human CD86 (NP_787058.4).
Preparation and Storage	Store at -20°C. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human tonsil using CD86 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using CD86 Antibody at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-CD86 antibody	This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript variants encoding different isoforms.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	332634950
NCBI GeneID	942
NCBI Accession #	NP_001193853.1
NCBI GenBank Nucleotide #	NM_001206924.1
UniProt Accession #	P42081; Q13655; Q6FHB1; Q6GTS4; Q7M4L5; A0N0P0; B7Z2F3; B7Z702; E7ETN5; E9PC27
Molecular Weight	Calculated MW: 12kDa/24kDa/31kDa/37kDa Observed MW: 65kDa
NCBI Official Full Name	T-lymphocyte activation antigen CD86 isoform 4
NCBI Official Synonym Full Names	CD86 molecule
NCBI Official Symbol	CD86
NCBI Official Synonym Symbols	B70; B7-2; B7.2; LAB72; CD28LG2
NCBI Protein Information	T-lymphocyte activation antigen CD86; BU63; FUN-1; CTLA-4 counter-receptor B7.2; B-lymphocyte activation antigen B7-2; CD86 antigen (CD28 antigen ligand 2, B7-2 antigen)
UniProt Protein Name	T-lymphocyte activation antigen CD86
UniProt Gene Name	CD86
UniProt Synonym Gene Names	CD28LG2
UniProt Entry Name	CD86_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd86-antibody-immunohistochemistry-western-blot-elisa/37119>