

Rabbit MAPK3-T202 Antibody | anti-MAPK3 antibody

Phospho-MAPK3-T202 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA37352	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MAPK3-T202; Phospho-MAPK3-T202 Polyclonal Antibody; Phospho-MAPK3-T202: ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK; anti-MAPK3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	379
Applicable Applications for anti-MAPK3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Route	Synthetic Peptide
Immunogen	A phospho specific peptide corresponding to residues surrounding T202 of human MAPK3
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic, nuclear staining using Phospho-MAPK3-T202 antibody.



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human breast carcinoma tissue, using Phospho-MAPK3-T202 antibody.



WB (Western Blot)

Western blot analysis of extracts of Jurkat cells, using Phospho-MAPK3-T202 antibody at 1:2000 dilution. Jurkat cells were treated by PMA/TPA (200nM) for 10 minutes. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% BSA. Detection: ECL Enhanced Kit. Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-MAPK3 antibody	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act in a signaling cascade that regulates various cellular processes such as proliferation, differentiation, and cell cycle progression in response to a variety of extracellular signals. This kinase is activated by upstream kinases, resulting in its translocation to the nucleus where it phosphorylates nuclear targets. Alternatively spliced transcript variants encoding different protein isoforms have been described.
Product Categories/Family for anti-MAPK3 antibody	Phosphorylation

NCBI AND UNIPROT INFORMATION

NCBI GI #	232066
NCBI GeneID	5595
UniProt Accession #	P27361; Q8NHX1; A8CZ58; B0LPG3
Molecular Weight	379
NCBI Official Full Name	Mitogen-activated protein kinase 3
NCBI Official Synonym Full Names	mitogen-activated protein kinase 3
NCBI Official Symbol	MAPK3
NCBI Official Synonym Symbols	ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK
NCBI Protein Information	mitogen-activated protein kinase 3; MAPK 1; MAP kinase isoform p44; insulin-stimulated MAP2 kinase; extracellular signal-related kinase 1; extracellular signal-regulated kinase 1; microtubule-associated protein 2 kinase
UniProt Protein Name	Mitogen-activated protein kinase 3
UniProt Gene Name	MAPK3
UniProt Synonym Gene Names	ERK1; PRKM3; MAP kinase 3; MAPK 3; ERK-1; p44-MAPK
UniProt Entry Name	MK03_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.