

Rabbit LIMK2-T505 Antibody | anti-LIMK2 antibody

Phospho-LIMK2-T505 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA37370	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

LIMK2-T505; Phospho-LIMK2-T505 Polyclonal Antibody; Phospho-LIMK2-T505: N/A; anti-LIMK2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	638
Applicable Applications for anti-LIMK2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Route	Synthetic Peptide
Immunogen	A phospho specific peptide corresponding to residues surrounding T505 of human LIMK2
Calculated Molecular Weight	70kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


IMAGES


IF (Immunofluorescence)

Immunofluorescence staining of methanol-fixed HeLa cells using Phospho-LIMK2-T505 antibody.


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human breast carcinoma using Phospho-LIMK2-T505 antibody.



WB (Western Blot)

Western blot analysis of extracts from HeLa cells, using Phospho-LIMK2-T505 antibody. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% BSA.

ADDITIONAL INFORMATION

Related Product Information for anti-LIMK2 antibody	There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. The protein encoded by this gene is phosphorylated and activated by ROCK, a downstream effector of Rho, and the encoded protein, in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity. It is thought that this pathway contributes to Rho-induced reorganization of the actin cytoskeleton. At least three transcript variants encoding different isoforms have been found for this gene.
Product Categories/Family for anti-LIMK2 antibody	Phosphorylation

NCBI AND UNIPROT INFORMATION

NCBI GI #	1708824
NCBI GeneID	3985
UniProt Accession #	P53671; Q7KZ80; Q7L3H5; Q96E10; Q99464; Q9UFU0; A8K6H5
Molecular Weight	638
NCBI Official Full Name	LIM domain kinase 2
NCBI Official Synonym Full Names	LIM domain kinase 2
NCBI Official Symbol	LIMK2
NCBI Protein Information	LIM domain kinase 2
UniProt Protein Name	LIM domain kinase 2
UniProt Gene Name	LIMK2
UniProt Synonym Gene Names	LIMK-2
UniProt Entry Name	LIMK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rabbit-human-mouse-rat-antibody/limk2-t505/37370>