

Rabbit PSMA7 Polyclonal Antibody | anti-PSMA7 antibody

PSMA7 Polyclonal Antibody

CATALOG # AAA37529	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
------------------------------	-----------------------	--	---



PRODUCT SYNONYMS

PSMA7; Polyclonal Antibody; PSMA7 Polyclonal Antibody; C6; HSPC; RC6-1; XAPC7; anti-PSMA7 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	248
Applicable Applications for anti-PSMA7 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human PSMA7
Calculated Molecular Weight	28kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



WB (Western Blot)
Western blot analysis of extracts of various cell lines, using PSMA7 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.

WB (Western Blot)
Western blot analysis of extracts of rat liver, using PSMA7 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.

ADDITIONAL INFORMATION

Related Product Information for anti-PSMA7 antibody	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. This gene encodes a member of the peptidase T1A family that functions as a 20S core alpha subunit. The encoded protein interacts with the hepatitis B virus X protein and plays a role in regulating hepatitis C virus internal ribosome entry site (IRES) activity, an activity essential for viral replication. The encoded protein also plays a role in the cellular stress response by regulating hypoxia-inducible factor-1alpha. A pseudogene of this gene is located on the long arm of chromosome 9.
Product Categories/Family for anti-PSMA7 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	12643540
NCBI GeneID	5688
UniProt Accession #	O14818 ; Q5JXJ2 ; Q9BR53 ; Q9H4K5 ; Q9UEU8 ; B2R515
Molecular Weight	248
NCBI Official Full Name	Proteasome subunit alpha type-7
NCBI Official Synonym Full Names	proteasome (prosome, macropain) subunit, alpha type, 7
NCBI Official Symbol	PSMA7
NCBI Official Synonym Symbols	C6; HSPC; RC6-1; XAPC7
NCBI Protein Information	proteasome subunit alpha type-7; proteasome subunit RC6-1; proteasome subunit XAPC7; proteasome subunit alpha 4
UniProt Protein Name	Proteasome subunit alpha type-7
UniProt Gene Name	PSMA7
UniProt Synonym Gene Names	HSPC
UniProt Entry Name	PSA7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/psma7-antibody-immunohistochemistry-western-blot-elisa/37529>