

Rabbit anti-Human, Mouse PSMC2 Polyclonal Antibody | anti-PSMC2 antibody

PSMC2 Polyclonal Antibody

CATALOG # AAA37535	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Immunofluorescence, Immunohistochemistry, Western Blot
------------------------------	-----------------------	-----------------------------------	---


PRODUCT SYNONYMS

PSMC2; Polyclonal Antibody; PSMC2 Polyclonal Antibody; S7; MSS1; Nbla10058; anti-PSMC2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	433
Applicable Applications for anti-PSMC2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human PSMC2
Calculated Molecular Weight	48kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.



IMAGES


IF (Immunofluorescence)

Immunofluorescence analysis of U2OS cells using PSMC2 antibody. Blue: DAPI for nuclear staining.


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse brain using PSMC2 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)
Western blot analysis of extracts of various cell lines, using PSMC2 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3s.

IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded mouse heart using PSMC2 antibody at dilution of 1:100 (40x lens).

ADDITIONAL INFORMATION

Related Product Information for anti-PSMC2 antibody	The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. This subunit has been shown to interact with several of the basal transcription factors so, in addition to participation in proteasome functions, this subunit may participate in the regulation of transcription. This subunit may also compete with PSMC3 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex. Alternative splicing results in multiple transcript variants encoding distinct isoforms.
Product Categories/Family for anti-PSMC2 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	547930
NCBI GeneID	5701
UniProt Accession #	P35998; Q3LIA5; Q9UDI3; A4D0Q1; B7Z5E2
Molecular Weight	433
NCBI Official Full Name	26S protease regulatory subunit 7
NCBI Official Synonym Full Names	proteasome (prosome, macropain) 26S subunit, ATPase, 2
NCBI Official Symbol	PSMC2
NCBI Official Synonym Symbols	S7; MSS1; Nbla10058
NCBI Protein Information	26S protease regulatory subunit 7; protease 26S subunit 7; proteasome 26S subunit ATPase 2; proteasome 26S subunit, ATPase, 2; putative protein product of Nbla10058; 26S proteasome AAA-ATPase subunit RPT1; mammalian suppressor of sgV-1 of yeast
UniProt Protein Name	26S protease regulatory subunit 7
UniProt Gene Name	PSMC2
UniProt Synonym Gene Names	MSS1
UniProt Entry Name	PRS7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/psmc2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/37535>