

Rabbit IL3RA Polyclonal Antibody | anti-IL3RA antibody

IL3RA Polyclonal Antibody

CATALOG # AAA37550	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

IL3RA; Polyclonal Antibody; IL3RA Polyclonal Antibody; CD123; IL3R; IL3RAY; IL3RX; IL3RY; hIL-3Ra; interleukin-3 receptor subunit alpha; anti-IL3RA antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Sequence Length	378
Applicable Applications for anti-IL3RA antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human IL3RA (NP_002174.1).
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffinembedded mouse kidney using IL3RA antibody (AAA37550) at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffinembedded rat lung using IL3RA antibody (AAA37550) at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of A-549 cells, using IL3RA antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-IL3RA antibody	The protein encoded by this gene is an interleukin 3 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL3 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL3. This gene and the gene encoding the colony stimulating factor 2 receptor alpha chain (CSF2RA) form a cytokine receptor gene cluster in a X-Y pseudoautosomal region on chromosomes X or Y. Alternatively spliced transcript variants encoding distinct isoforms have been found.
Product Categories/Family for anti-IL3RA antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	417184
NCBI GeneID	3563
UniProt Accession #	P26951; Q5HYQ7; Q5HYQ8; Q9UEH7; A8K3F3; B9VI81
Molecular Weight	Observed MW: 70kDa Calculated MW: 34kDa/43kDa
NCBI Official Full Name	Interleukin-3 receptor subunit alpha
NCBI Official Synonym Full Names	interleukin 3 receptor, alpha (low affinity)
NCBI Official Symbol	IL3RA
NCBI Official Synonym Symbols	IL3R; CD123; IL3RX; IL3RY; IL3RAY; hIL-3Ra
NCBI Protein Information	interleukin-3 receptor subunit alpha; IL-3RA; IL-3R-alpha; CD123 antigen; IL-3R subunit alpha; IL-3 receptor subunit alpha; IL-3 receptor alpha SP2 isoform
UniProt Protein Name	Interleukin-3 receptor subunit alpha
UniProt Gene Name	IL3RA
UniProt Synonym Gene Names	IL3R; IL-3 receptor subunit alpha; IL-3R subunit alpha; IL-3R-alpha; IL-3RA
UniProt Entry Name	IL3RA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

