

Rabbit WISP2 Polyclonal Antibody | anti-WISP2 antibody

WISP2

CATALOG # AAA37625	HOST Rabbit	APPLICATIONS Immunohistochemistry, Western Blot
------------------------------	-----------------------	---


PRODUCT SYNONYMS

WISP2; Polyclonal Antibody; CCN5; CT58; CTGF-L; WNT1-inducible-sign aling pathway protein 2; anti-WISP2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.09% Sodium azide,50% glycerol,pH7.3.
Sequence Length	250
Applicable Applications for anti-WISP2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human WISP2 (NP_003872.1).
Species Reactivity/Cross-Reactivity	Human, Mouse, Rat
Category	Primary Antibody
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles. Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

IMAGES




IHC (Immunohistochemisry)

Immunohistochemistry of paraffin-embedded human colon using WISP2 Rabbit pAb (AAA37625) at dilution of 1:100 (40x lens).



IHC (Immunohiostchemistry)

Immunohistochemistry of paraffin-embedded rat testis using WISP2 Rabbit pAb (AAA37625) at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using WISP2 antibody (AAA37625) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-WISP2 antibody	This gene encodes a member of the WNT1 inducible signaling pathway (WISP) protein subfamily, which belongs to the connective tissue growth factor (CTGF) family. WNT1 is a member of a family of cysteine-rich, glycosylated signaling proteins that mediate diverse developmental processes. The CTGF family members are characterized by four conserved cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C module, thrombospondin domain and C-terminal cystine knot-like (CT) domain. The encoded protein lacks the CT domain which is implicated in dimerization and heparin binding. It is 72% identical to the mouse protein at the amino acid level. This gene may be downstream in the WNT1 signaling pathway that is relevant to malignant transformation. Its expression in colon tumors is reduced while the other two WISP members are overexpressed in colon tumors. It is expressed at high levels in bone tissue, and may play an important role in modulating bone turnover.
Product Categories/Family for anti-WISP2 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	24211561
NCBI GeneID	8839
UniProt Accession #	O76076 ; Q6PEG3 ; B2R9N4 ; E1P612
Molecular Weight	Observed: 27kDa Calculated: 22kDa/26kDa
NCBI Official Full Name	WNT1-inducible-signaling pathway protein 2
NCBI Official Synonym Full Names	WNT1 inducible signaling pathway protein 2
NCBI Official Symbol	WISP2
NCBI Official Synonym Symbols	CCN5; CT58; CTGF-L
NCBI Protein Information	WNT1-inducible-signaling pathway protein 2; WISP-2; CCN family member 5; wnt-1 signaling pathway protein 2; connective tissue growth factor-like protein; connective tissue growth factor-related protein 58
UniProt Protein Name	WNT1-inducible-signaling pathway protein 2
UniProt Gene Name	WISP2
UniProt Synonym Gene Names	CCN5; CT58; CTGFL; WISP-2; CTGF-L
UniProt Entry Name	WISP2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/wisp2-antibody-immunohistochemistry-western-blot-elisa/37625>