

Monocyte Chemotactic Protein-1 Active Protein | rMCP 1 active protein

Recombinant Rat Monocyte Chemotactic Protein-1 (CCL2)

CATALOG #
AAA38086

HOST
E Coli

PRODUCT SYNONYMS

Monocyte Chemotactic Protein-1; Recombinant Rat Monocyte Chemotactic Protein-1 (CCL2); MCP 1 Rat; Monocyte Chemotactic Protein-1 Rat Recombinant (CCL2); Small inducible cytokine A2; CCL2; Monocyte chemotactic protein 1; MCP-1; Monocyte chemoattractant protein 1; Monocyte chemotactic and activating factor; MCAF; Monocyte secretory protein JE; HC11; chemokine (C-C motif) ligand 2; MCP1; SCYA2; GDCF-2; SMC-CF; HSMCR30; MGC9434; GDCF-2 HC11; Immediate-early serum-responsive JE protein; rMCP 1 active protein

PRODUCT SPECIFICATIONS

Host	E Coli
Purity/Purification	Greater than 98.0% as determined by SDS-PAGE.
Form/Format	The protein was lyophilized from a concentrated (1mg/ml) sterile solution containing no additives. Sterile Filtered White lyophilized (freeze-dried) powder.
Sequence	QPDVNAPLT CCYSFTGKMI PMSRLNYKR ITSSRCPKEA VVFVTKLKRE ICADPNKEWV QKYIRKLDQN QVRSETTVFY KIASLRTSA PLNVNLTHKS EANASTLFST TTSSTSVEVT SMTE
Sequence Length	148
Solubility	It is recommended to reconstitute the lyophilized Monocyte Chemotactic Protein-1 in sterile 18M Omega -cm H2O not less than 100 ug/ml, which can then be further diluted to other aqueous solutions.
Biological Activity	ED50 =1-10ng/ml corresponding to a Specific Activity of 100,000-1,000,000IU/mg. The biological activity was determined by measuring the dose dependent chemotaxis with human THP-1 cells. The optimal concentration should be determined for each specific application by an initial dose-response assay.
Preparation and Storage	Lyophilized MCP-1 although stable at room temperature for 3 weeks, should be stored desiccated below -18 degree C. Upon reconstitution CCL2 should be stored at 4 degree C between 2-7 days and for future use below -18 degree C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

ADDITIONAL INFORMATION

Related Product Information for rMCP 1 active protein	Description: Monocyte Chemotactic Protein-1 Rat Recombinant produced in E Coli is a non-glycosylated, Polypeptide chain containing 125 amino acids and having a molecular mass of 14.1 kDa. The MCP-1 is purified by proprietary chromatographic techniques. Introduction: Chemokine (C-C motif) ligand 2 (CCL2) is a small cytokine belonging to the CC chemokine family that is also known as monocyte chemotactic protein-1 (MCP-1). It is found at the site of tooth eruption and bone degradation. In the bone, CCL2 is expressed by mature osteoclasts and osteoblasts and is under the control of nuclear factor κB (NFκB). CCL2 recruits immune cells, such as monocytes, to sites of tissue injury and infection. This chemokine is produced as a protein precursor containing signal peptide of 23 amino acids and a mature peptide of 76 amino acids. It is a monomeric polypeptide, with a molecular weight of approximately 13kDa. As with many other CC chemokines, CCL2 is located on chromosome 17 in humans. The cell surface receptors that bind CCL2 are CCR2 and CCR5.
Product Categories/Family for rMCP 1 active protein	Chemokines; MCP

NCBI AND UNIPROT INFORMATION

NCBI GI #	13928714
NCBI GeneID	24770
NCBI Accession #	NP_113718.1
NCBI GenBank Nucleotide #	NM_031530.1
UniProt Accession #	P14844; Q549R5
Molecular Weight	16,460 Da
NCBI Official Full Name	C-C motif chemokine 2
NCBI Official Synonym Full Names	chemokine (C-C motif) ligand 2
NCBI Official Symbol	Ccl2
NCBI Official Synonym Symbols	MCP-1; Scya2; Sigje
NCBI Protein Information	C-C motif chemokine 2; immediate-early serum-responsive JE protein; immediate-early serum-responsive protein JE; monocyte chemoattractant protein 1; monocyte chemoattractant protein-1; monocyte chemotactic protein 1; small inducible cytokine A2; small inducible gene JE; small-inducible cytokine A2
UniProt Protein Name	C-C motif chemokine 2
UniProt Gene Name	Ccl2
UniProt Synonym Gene Names	Je; Mcp1; Scya2; MCP-1
UniProt Entry Name	CCL2_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/e-coli-rmcp-1-active-protein/monocyte-chemotactic-protein-1/38086>