

Granulocyte Macrophage-Colony Stimulating Factor Active Protein | mGM CSF active protein

Recombinant Mouse Granulocyte Macrophage-Colony Stimulating Factor

CATALOG #
AAA38116

PRODUCT SYNONYMS

Granulocyte Macrophage-Colony Stimulating Factor; Recombinant Mouse Granulocyte Macrophage-Colony Stimulating Factor; GM-CSF Mouse; Granulocyte Macrophage-Colony Stimulating Factor Mouse Recombinant; CSF-2; MGI-1GM; GM-CSF; Pluripoietin-alpha; Molgramostin; Sargramostim; mGM-CSF active protein

PRODUCT SPECIFICATIONS

Purity/Purification	Greater than 98.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.
Form/Format	GM-CSF Mouse was lyophilized with no additives. Sterile Filtered White lyophilized (freeze-dried) powder.
Sequence	Thesequenc eofthefirs tfiveN-ter minalamino acidswasde terminedan dwasfoundt obeMet-Ala -Pro-Thr-A rg
Sequence Length	141
Source	E Coli
Solubility	It is recommended to reconstitute the lyophilized Granulocyte Macrophage Colony Stimulating Factor in sterile 20mM AcOH (acetic Acid) not less than 100 ug/ml, which can then be further diluted to other aqueous solutions.
Protein Content	GM-CSF quantitation was carried out by two independent methods 1. UV spectroscopy at 280 nm using the absorbency value of 0.765 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the PC GENE computer analysis program of protein sequences (IntelliGenetics). 2. Analysis by RP-HPLC, using a calibrated solution of GM-CSF as a Reference Standard.
Biological Activity	The ED ₅₀ as determined by the dose-dependant stimulation of the proliferation of murine FDC-P1 cell line is < 0.2 ng/ml, corresponding to a Specific Activity of 5 x 10 ⁶ IU/mg.
Preparation and Storage	Lyophilized Granulocyte Macrophage Colony Stimulating Factor although stable at room temperature for 3 weeks, should be stored desiccated below -18 degree C. Upon reconstitution GM-CSF should be stored at 4 degree C between 2-7 days and for future use below -18 degree C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

ADDITIONAL INFORMATION

Related Product Information for mGM CSF active protein	Description: Granulocyte Macrophage Colony Stimulating Factor Mouse Recombinant produced in E Coli is a single, non-glycosylated, polypeptide chain containing 125 amino acids and having a molecular mass of 14285.35 Dalton. GM-CSF Mouse is purified by proprietary chromatographic techniques. Introduction: GM-CSF is a cytokine that controls the production, differentiation, and function of granulocytes and macrophages. The active form of the protein is found extracellularly as a homodimer. This gene has been localized to a cluster of related genes at chromosome region 5q31, which is known to be associated with interstitial deletions in the 5q- syndrome and acute myelogenous leukemia. Other genes in the cluster include those encoding interleukins 4, 5, and 13. GM-CSF stimulates the growth and differentiation of hematopoietic precursor cells from various lineages, including granulocytes, macrophages, eosinophils and erythrocytes.
Product Categories/Family for mGM CSF active protein	Cytokines and Growth Factors; Growth Factors; Colony Stimulating Factor

NCBI AND UNIPROT INFORMATION

NCBI GI #	110625658
NCBI GeneID	12981
NCBI Accession #	NP_034099.2
NCBI GenBank Nucleotide #	NM_009969.4
UniProt Accession #	P01587
Molecular Weight	16,091 Da
NCBI Official Full Name	granulocyte-macrophage colony-stimulating factor
NCBI Official Synonym Full Names	colony stimulating factor 2 (granulocyte-macrophage)
NCBI Official Symbol	Csf2
NCBI Official Synonym Symbols	Csfgm; GMCSF; Gm-CSf; MGI-IGM
NCBI Protein Information	granulocyte-macrophage colony-stimulating factor; CSF; granulocyte-macrophage colony stimulating factor 2; put. GM-CSF
UniProt Protein Name	Granulocyte-macrophage colony-stimulating factor
UniProt Gene Name	Csf2
UniProt Synonym Gene Names	Csfgm; GM-CSF; CSF
UniProt Entry Name	CSF2_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mgm-csf-active-protein/granulocyte-macrophage-colony-stimulating-factor/38116>