

Interleukin-15 Active Protein | rIL 15 active protein

Recombinant Rat Interleukin-15

CATALOG #
AAA38169

HOST
E Coli

PRODUCT SYNONYMS

Interleukin-15; Recombinant Rat Interleukin-15; IL 15 Rat; Interleukin-15 Rat Recombinant; IL-15; MGC9721; rIL 15 active protein

PRODUCT SPECIFICATIONS

Host	E Coli
Purity/Purification	Greater than 98.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.
Form/Format	The protein was lyophilized from 10mM Tris, pH-8.5.
Sequence	Thesequenc eofthefirs tfiveN-ter minalamino acidswasde terminedan dwasfoundt obeMet-Asn -Trp-Ile-A sp
Sequence Length	162
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder).
Solubility	It is recommended to reconstitute the lyophilized Interleukin-15 in sterile 18M Omega -cm H2O not less than 100 ug/ml, which can then be further diluted to other aqueous solutions.
Biological Activity	The ED50 as determined by the dose-dependant stimulation of the proliferation of CTLL-2 was found to be < 10 ng/ml, corresponding to a Specific Activity of 1×10^5 IU/mg.
Preparation and Storage	Lyophilized Interleukin-15 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL15 should be stored at 4° between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

ADDITIONAL INFORMATION

Related Product Information for rIL 15 active protein	Description: Interleukin-15 Rat Recombinant produced in E Coli is a single, non-glycosylated polypeptide chain containing 115 amino acids and having a molecular mass of 13533 Dalton. The IL-15 is purified by proprietary chromatographic techniques. Introduction: The protein encoded by this gene is a cytokine that regulates T and natural killer cell activation and proliferation. This cytokine and interleukine 2 share many biological activities. They are found to bind common hematopoietin receptor subunits, and may compete for the same receptor, and thus negatively regulate each other's activity. The number of CD8+ memory cells is shown to be controlled by a balance between this cytokine and IL2. This cytokine induces the activation of JAK kinases, as well as the phosphorylation and activation of transcription activators STAT3, STAT5, and STAT6. Studies of the mouse counterpart suggested that this cytokine may increase the expression of apoptosis inhibitor BCL2L1/BCL-x(L), possibly through the transcription activation activity of STAT6, and thus prevent apoptosis. Two alternatively spliced transcript variants of this gene encoding the same protein have been reported.
Product Categories/Family for rIL 15 active protein	Cytokines and Growth Factors ; Cytokines ; Interleukin

NCBI AND UNIPROT INFORMATION

NCBI GI #	6981092
NCBI GeneID	25670
NCBI Accession #	NP_037261.1
NCBI GenBank Nucleotide #	NM_013129.2
UniProt Accession #	P97604; O54847
NCBI Official Full Name	interleukin-15
NCBI Official Synonym Full Names	interleukin 15
NCBI Official Symbol	IL15
NCBI Protein Information	interleukin-15; IL-15
UniProt Protein Name	Interleukin-15
UniProt Gene Name	IL15
UniProt Synonym Gene Names	IL-15
UniProt Entry Name	IL15_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/e-coli-ril-15-active-protein/interleukin-15/38169>