

HIV-1 gp41 Long Recombinant Protein

Recombinant HIV-1 gp41 Long, Horseradish peroxidase Labeled

CATALOG #
AAA38334

HOST
Escherichia Coli.

PRODUCT SYNONYMS

HIV-1 gp41 Long; Recombinant HIV-1 gp41 Long; Horseradish peroxidase Labeled; HRP; HIV-1 gp41 Long Recombinant; HRP Labeled; HIV-1 gp41 L-HRP; HIV-1 gp41 Long recombinant protein

PRODUCT SPECIFICATIONS

Host	Escherichia Coli.
Specificity	Immunoreactive with HIV positive serum.
Purity/Purification	Greater than 95.0% as determined by HPLC and SDS-PAGE.
Form/Format	8M urea, 20mM Tris-HCl pH 8.0, 10mM b-mercaptoethanol.
Physical Appearance	Sterile Filtered clear solution (1mg/1mL)
Preparation and Storage	HIV-1 gp41 Long HRP although stable at 4°C for 1 week, should be stored below -18°C. Please prevent freeze thaw cycles.

ADDITIONAL INFORMATION

Related Product Information for HIV-1 gp41 Long recombinant protein	Description: The E Coli derived HRP Labeled protein contains HIV immunodominant regions from HIV-I gp41 and is fused to beta-galactosidase (114 kDa) at the N-terminus. Introduction: Human immunodeficiency virus (HIV) is a retrovirus that can lead to a condition in which the immune system begins to fail, leading to opportunistic infections. HIV primarily infects vital cells in the human immune system such as helper T cells (specifically CD4+ T cells), macrophages and dendritic cells. HIV infection leads to low levels of CD4+ T cells through three main mechanisms: firstly, direct viral killing of infected cells; secondly, increased rates of apoptosis in infected cells; and thirdly, killing of infected CD4+ T cells by CD8 cytotoxic lymphocytes that recognize infected cells. When CD4+ T cell numbers decline below a critical level, cell-mediated immunity is lost, and the body becomes progressively more susceptible to opportunistic infections. HIV was classified as a member of the genus Lentivirus, part of the family of Retroviridae. Lentiviruses have many common morphologies and biological properties. Many species are infected by lentiviruses, which are characteristically responsible for long-duration illnesses with a long incubation period. Lentiviruses are transmitted as single-stranded, positive-sense, enveloped RNA viruses. Upon entry of the target cell, the viral RNA genome is converted to double-stranded DNA by a virally encoded reverse transcriptase that is present in the virus particle. This viral DNA is then integrated into the cellular DNA by a virally encoded integrase so that the genome can be transcribed. Once the virus has infected the cell, two pathways are possible: either the virus becomes latent and the infected cell continues to function, or the virus becomes active and replicates, and a large number of virus particles are liberated that can then infect other cells.
Product Categories/Family for HIV-1 gp41 Long recombinant protein	Viral Antigens; HUMAN IMMUNODEFICIENCY VIRUS; Viral Antigens; HIV

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.