

Rabbit Caspase-12 Polyclonal Antibody | anti-Casp12 antibody

Caspase-12 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA39494	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

Caspase-12; Polyclonal Antibody; Caspase-12 Antibody; CASP-12; caspase 12; anti-Casp12 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Caspase-12 Antibody is affinity chromatography purified via peptide column.
Form/Format	Liquid
Sequence Length	419
Applicable Applications for anti-Casp12 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Conjugate	Unconjugated
Immunogen	Caspase-12 antibody was raised against a 17 amino acid peptide near the center of murine Caspase-12.
Buffer	Caspase-12 Antibody is supplied in PBS containing 0.02% sodium azide.
Preparation and Storage	Caspase-12 antibody can be stored at 4 degree C for three months and -20 degree C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.


IMAGES


IF (Immunofluorescence)

Immunofluorescence of Caspase-12 in Mouse Liver cells with Caspase-12 antibody at 10 µg/mL.


IHC (Immunohistochemistry)

Immunohistochemical staining of mouse liver tissue using caspase-12 antibody at 2 µg/mL.



WB (Western Blot)

Western blot analysis of Caspase-12 in (A) human spleen and (B) mouse spleen tissue lysate with Caspase-12 antibody at 1 µg/mL.

ADDITIONAL INFORMATION

Related Product Information for anti-Casp12 antibody

Caspase-12 Antibody: Three distinct signaling pathways lead to programmed cell death (apoptosis). The death receptor and mitochondrion pathways are the main, in which the key apoptotic proteases caspase-8 and caspase-9, respectively, are involved. The endoplasmic reticulum (ER) stress is the third apoptotic pathway and caspase-12 is involved. Caspase-12 is localized to the ER but not to cytoplasm or mitochondrion. Caspase-12 is activated by ER stress, including disruption of ER calcium homeostasis, and mediates ER stress-induced apoptosis. Caspase-12 is co-localized to the ER with several proteins that are involved in Alzheimer's disease including gamma-secretase presenilin and beta-amyloid precursor protein (APP). Caspase-12 mediates cytotoxicity induced by amyloid-beta. Caspase-12 is ubiquitously expressed in mouse tissues.

NCBI AND UNIPROT INFORMATION

NCBI GI #	2094806
NCBI GeneID	12364
UniProt Accession #	O08736; Q3TT82
Molecular Weight	47,854 Da
NCBI Official Full Name	caspase-12
NCBI Official Synonym Full Names	caspase 12
NCBI Official Symbol	Casp12
NCBI Protein Information	caspase-12; CASP-12
UniProt Protein Name	Caspase-12
UniProt Gene Name	Casp12
UniProt Synonym Gene Names	CASP-12
UniProt Entry Name	CASPC_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/caspase-12-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/39494>