

Mouse anti-Human PD-1 Monoclonal Antibody | anti-PDCD1 antibody

PD-1 Antibody [4D6]

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA40879	Mouse	Human	Immunofluorescence, Immunocytochemistry, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

PD-1; Monoclonal Antibody; PD-1 Antibody [4D6]; PD-1 Antibody; PD1; CD279; SLEB2; hPD-1; hPD-I; hSLE1; Programmed cell death protein 1; Protein PD-1; PDCD1; PDCD-1; anti-PDCD1 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Clone Number	4D6
Purity/Purification	Protein A purified IgG1.
Form/Format	Liquid
Concentration	1 mg/mL (varies by lot)
Sequence Length	288
Applicable Applications for anti-PDCD1 antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Conjugate	Unconjugated
Immunogen	PD-1 antibody was raised against the extracellular domain of human PD-1.
Buffer	PD-1 Antibody is supplied in PBS containing 0.02% sodium azide.
Species	Human
Preparation and Storage	Store at 4 degree C for three months and -20 degree C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles Antibodies should not be exposed to prolonged high temperatures.

IMAGES



WB (Western Blot)

Western blot analysis of PD-1 in overexpressing 293 cells using antibody at 1, 0.5, and 0.25 µg/mL, respectively.




IF (Immunofluorescence)

Immunofluorescence of PD-1 in transfected 293 cells with PD-1 antibody at 5 µg/mL. Lower left: Immunofluorescence in transfected 293 cells with control mouse IgG antibody at 5 µg/mL. Red: PD1 Antibody [4D6] Blue: DAPI staining




ICC (Immunocytochemistry)

Immunocytochemistry of PD-1 in transfected 293 cells with PD-1 antibody at 5 μg/mL. Lower left: Immunocytochemistry in transfected 293 cells with control mouse IgG antibody at 5 μg/mL.



IHC (Immunohistochemistry)

Immunohistochemistry of PD-1 in (A) human tonsil tissue, (B) human lymph node tissue, and (C) human spleen tissue with PD-1 antibody at 5 μg/mL. (D) Immunohistochemistry in human tonsil tissue with control mouse IgG staining at 5 μg/mL.



WB (Western Blot)

Western blot analysis of PD-1 in transfected 293 cell lysate with PD-1 antibody at (A) 0.25 and (B) 0.5 μg/mL.

ADDITIONAL INFORMATION

Related Product Information for anti-PDCD1 antibody	Cell-mediated immune responses are initiated by T lymphocytes that are themselves stimulated by cognate peptides bound to MHC molecules on antigen-presenting cells (APC). T-cell activation is generally self-limited as activated T cells express receptors such as PD-1 (also known as PDCD-1) that mediate inhibitory signals from the APC. PD-1 can bind two different but related ligands, PDL-1 and PDL-2. Upon binding to either of these ligands, signals generated by PD-1 inhibit the activation of the immune response in the absence of "danger signals" such as LPS or other molecules associated with bacteria or other pathogens. Evidence for this is seen in PD-1-null mice who exhibit hyperactivated immune systems and autoimmune diseases. PD-1 is thus one of a growing number of immune checkpoint proteins.
Product Categories/Family for anti-PDCD1 antibody	Apoptosis
References	1. Holling TM, Schooten E, and van Den Elsing PJ. Function and regulation of MHC class II molecules in T-lymphocytes: of mice and men. Hum. Immunol. 2004; 65:282-90. 2. Ishida Y, Agata Y, Shibahara K, et al. Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. EMBO J. 1992; 11:3887-95. 3. Zhong X, Bai C, Gao W, et al. Suppression of expression and function of negative immune regulator PD-1 by certain pattern recognition and cytokine receptor signals associated with immune system danger. Int. Immunol. 2004; 16:1181-8. 4. Nishimura H, Nose M, Hiai H, et al. Development of lupus-like autoimmune diseases by the disruption of the PD-1 gene encoding an ITIM motif-carrying immunoreceptor. Immunity 1999; 11:141-51.

NCBI AND UNIPROT INFORMATION

NCBI GI #	145559515
NCBI GeneID	5133
UniProt Accession #	Q15116 ; O00517 ; Q8IX89
Molecular Weight	Predicted: 32 kDa Observed: 38 kDa
NCBI Official Full Name	Programmed cell death protein 1
NCBI Official Synonym Full Names	programmed cell death 1
NCBI Official Symbol	PDCD1
NCBI Official Synonym Symbols	PD1; PD-1; CD279; SLEB2; hPD-1; hPD-I; hSLE1
NCBI Protein Information	programmed cell death protein 1
UniProt Protein Name	Programmed cell death protein 1
UniProt Gene Name	PDCD1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/pd-1-antibody-clone-4d6-immunofluorescence-immunocytochemistry-immunohistochemistry-western-blot-elisa/40879>