

Human High Mobility Group Protein 17, HMG-17 ELISA Kit | HMG-17 elisa kit

Human High Mobility Group Protein 17, HMG-17 ELISA Kit

CATALOG #
AAA41258

REACTIVITY
Human


PRODUCT SYNONYMS

High Mobility Group Protein 17; HMG-17; Human High Mobility Group Protein 17; HMG-17 ELISA Kit; HMG-17 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	91
Samples	Serum, Plasma, Cell Culture Supernates, Cell Lysates, Tissue Homogenates
Assay Type	Quantitative Sandwich
Detection Range	0.1ng/ml-40ng/ml
Sensitivity	0.053ng/ml
Intra-assay Precision	Intra-Assay Precision (Precision within an assay) Three samples of known concentration were tested on one plate to assess intra-assay precision. Intra-Assay: CV<8%
Inter-assay Precision	Inter-Assay Precision (Precision between assays) Three samples of known concentration were tested in separate assays to assess inter-assay precision. CV(%) = SD/mean x 100. Inter-Assay: CV<10%
Preparation and Storage	Store at 2 to 8 degree C for 6 months.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HMG-17 elisa kit

Intended Uses: This sandwich kit is for the accurate quantitative detection of human Hexokinase-2 (also known as HK2) in serum, plasma, cell culture supernates, cell lysates, tissue homogenates.

Principle of the Assay: This kit is an Enzyme-Linked Immunosorbent Assay (ELISA). The plate has been pre-coated with human HK2 antibody. HK2 present in the sample is added and binds to antibodies coated on the wells. And then biotinylated human HK2 Antibody is added and binds to HK2 in the sample. Then Streptavidin-HRP is added and binds to the Biotinylated HK2 antibody. After incubation unbound Streptavidin-HRP is washed away during a washing step. Substrate solution is then added and color develops in proportion to the amount of human HK2. The reaction is terminated by addition of acidic stop solution and absorbance is measured at 450 nm.

NCBI AND UNIPROT INFORMATION

NCBI GI #	148234670
NCBI GeneID	398020
NCBI Accession #	NP_001081730.1
NCBI GenBank Nucleotide #	NM_001088261.1
NCBI Official Full Name	high mobility group nucleosomal binding domain 2 S homeolog
NCBI Official Synonym Full Names	high mobility group nucleosomal binding domain 2 S homeolog
NCBI Official Symbol	hmgn2.S
NCBI Official Synonym Symbols	hmg-17
NCBI Protein Information	high mobility group nucleosomal binding domain 2 S homeolog

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/high-mobility-group-protein-17-hmg-17/41258>