

## Mouse Vascular cell adhesion molecule 1, sVCAM-1 ELISA Kit | sVCAM-1 elisa kit

Mouse soluble Vascular cell adhesion molecule 1, sVCAM-1 ELISA Kit

CATALOG #  
**AAA41429**

REACTIVITY  
**Mouse**


### PRODUCT SYNONYMS

Vascular cell adhesion molecule 1; sVCAM-1; Mouse soluble Vascular cell adhesion molecule 1; sVCAM-1 ELISA Kit; sVCAM-1 elisa kit

### PRODUCT SPECIFICATIONS

|                         |                                                                          |
|-------------------------|--------------------------------------------------------------------------|
| Reactivity              | Mouse                                                                    |
| Samples                 | Serum, plasma, cell culture supernates, cell lysates, tissue homogenates |
| Assay Type              | Quantitative Sandwich                                                    |
| Detection Range         | 0.05ng/ml - 15ng/ml                                                      |
| Sensitivity             | 0.022ng/ml                                                               |
| Intra-Assay             | CV<8%                                                                    |
| Inter-Assay             | CV<10%                                                                   |
| Preparation and Storage | Store at 2 to 8 degree C for 6 months.                                   |

### IMAGES



Standard Curve (Sample)

### ADDITIONAL INFORMATION

#### Related Product Information for sVCAM-1 elisa kit

**Intended Uses:** This sandwich kit is for the accurate quantitative detection of Mouse Soluble Vascular Cell Adhesion Molecule 1 (also known as sVCAM-1) in serum, plasma, cell culture supernates, cell lysates, tissue homogenates.

**Principles of the Assay:** This kit is an Enzyme-Linked Immunosorbent Assay (ELISA). The plate has been pre-coated with Mouse sVCAM-1 antibody. sVCAM-1 present in the sample is added and binds to antibodies coated on the wells. And then biotinylated Mouse sVCAM-1 Antibody is added and binds to sVCAM-1 in the sample. Then Streptavidin-HRP is added and binds to the Biotinylated sVCAM-1 antibody. After incubation unbound Streptavidin-HRP is washed away during a washing step. Substrate solution is then added and color develops in proportion to the amount of Mouse sVCAM-1. The reaction is terminated by addition of acidic stop solution and absorbance is measured at 450 nm.

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | [contact@aaabiotech.com](mailto:contact@aaabiotech.com)

Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/vascular-cell-adhesion-molecule-1-svcam-1/41429>