

## Human lactoperoxidase (LPO) ELISA Kit | LPO elisa kit

Human lactoperoxidase (LPO) ELISA Kit

CATALOG #  
**AAA44402**

REACTIVITY  
**Human**


### PRODUCT SYNONYMS

lactoperoxidase (LPO); Human lactoperoxidase (LPO) ELISA Kit; LPO elisa kit

### PRODUCT SPECIFICATIONS

|                         |                                                                          |
|-------------------------|--------------------------------------------------------------------------|
| Reactivity              | Human                                                                    |
| Samples                 | Serum, plasma, cell culture supernates, cell lysates, tissue homogenates |
| Assay Type              | Quantitative Sandwich                                                    |
| Detection Range         | 0.05ng/ml - 9ng/ml                                                       |
| Sensitivity             | 0.023ng/ml                                                               |
| Intra-Assay             | CV<8%                                                                    |
| Inter-Assay             | CV<10%                                                                   |
| Preparation and Storage | Store at 2 to 8 degree C for 6 months.                                   |

### IMAGES



Standard Curve (Sample)

### ADDITIONAL INFORMATION

#### Related Product Information for LPO elisa kit

**Intended Uses:** This sandwich kit is for the accurate quantitative detection of human Lactoperoxidase (also known as LPO) in serum, plasma, cell culture supernates, cell lysates, tissue homogenates.

**Principles of the Assay:** This kit is an Enzyme-Linked Immunosorbent Assay (ELISA). The plate has been pre-coated with human LPO antibody. LPO present in the sample is added and binds to antibodies coated on the wells. And then biotinylated human LPO Antibody is added and binds to LPO in the sample. Then Streptavidin-HRP is added and binds to the Biotinylated LPO antibody. After incubation unbound Streptavidin-HRP is washed away during a washing step. Substrate solution is then added and color develops in proportion to the amount of human LPO. The reaction is terminated by addition of acidic stop solution and absorbance is measured at 450 nm.

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | [contact@aaabiotech.com](mailto:contact@aaabiotech.com)  
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/lactoperoxidase-lpo/44402>