

Human sCD40L ELISA Kit | CD40LG elisa kit

Human sCD40L PicoKine ELISA Kit

CATALOG #
AAA45308

REACTIVITY
Human


PRODUCT SYNONYMS

sCD40L; Human sCD40L PicoKine ELISA Kit; CD40 ligand; CD154; CD40 antigen ligand; CD40-L; CD40L_HUMAN; CD40LG; gp39; HIGM1; IGM; IMD3; soluble form; T B cell activating molecule; T BAM; T cell antigen Gp39; T-cell antigen Gp39; TNF related activation protein; TNF-related activation protein; TNFSF5; TrAP; Tumor necrosis factor (ligand) superfamily member 5; Tumor necrosis factor (ligand) superfamily; member 5 (hyper-IgM syndrome); Tumor necrosis factor ligand superfamily member 5; CD40LG elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	Natural and recombinant human CD40L
Sequence Length	261
Samples	Cell culture supernates, serum and plasma (heparin, EDTA)
Assay Type	Sandwich
Detection Range	62.5pg/ml-4000pg/ml
Sensitivity	< 15pg/ml
Intra-assay Precision	Intra-Assay Precision (Precision within an assay) Three samples of known concentration were tested on one plate to assess intra-assay precision.
Inter-assay Precision	Inter-Assay Precision (Precision between assays) Three samples of known concentration were tested in separate assays to assess inter-assay precision.
Preparation and Storage	Store at 4 degree C for 6 months, at -20 degree C for 12 months. Avoid multiple freeze-thaw cycles (Shipped with wet ice.)

IMAGES



Standard Curve (Sample)

Human sCD40L PicoKine ELISA Kit standard curve

ADDITIONAL INFORMATION

Related Product Information for CD40LG elisa kit

Principle of the assay: human CD40L ELISA Kit was based on standard sandwich enzyme-linked immune-sorbent assay technology. A monoclonal antibody from mouse specific for CD40L has been precoated onto 96-well plates. Standards(E.coli, E108-L261) and test samples are added to the wells, a biotinylated detection polyclonal antibody from goat specific for CD40L is added subsequently and then followed by washing with PBS or TBS buffer. Avidin-Biotin-Peroxidase Complex was added and unbound conjugates were washed away with PBS or TBS buffer. HRP substrate TMB was used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the human CD40L amount of sample captured in plate.

Background: CD40 ligand(CD40L) is a type II membrane protein of 261 amino acids on activated T cells that induces B cell proliferation and immunoglobulin secretion. It has homology with tumour necrosis factor-alpha and -beta, and has important functions in B-cell activation and differentiation. Human CD40L with 5 exons, is mapped to the proximal region of the mouse X chromosome on Xq26.3-27.1, and can be detected on T cells but is absent from B cells and monocytes. Since CD40L is expressed on platelets and released from them on activation, its predictive value as a marker for clinical outcome and the therapeutic effect of inhibition of glycoprotein IIb /IIIa receptor in patients with acute coronary syndromes was investigated. The soluble CD40L may be involved in the process of restenosis and that it exerts its effect by triggering a complex group of inflammatory reactions on endothelial and mononuclear cells. CD40L plays a central role in the pathophysiology of acute coronary syndromes, and has a role in the pathogenesis of coronary artery lesions.

NCBI AND UNIPROT INFORMATION

NCBI GI #	231718
NCBI GeneID	959
UniProt Accession #	P29965
Molecular Weight	29,274 Da
NCBI Official Full Name	CD40 ligand
NCBI Official Synonym Full Names	CD40 ligand
NCBI Official Symbol	CD40LG
NCBI Official Synonym Symbols	IGM; IMD3; TRAP; gp39; CD154; CD40L; HIGM1; T-BAM; TNFSF5; hCD40L
NCBI Protein Information	CD40 ligand; CD40-L; CD40 antigen ligand; T-cell antigen Gp39; T-B cell-activating molecule; TNF-related activation protein; tumor necrosis factor (ligand) superfamily member 5
UniProt Protein Name	CD40 ligand
UniProt Gene Name	CD40LG
UniProt Synonym Gene Names	CD40L; TNFSF5; TRAP; CD40-L; TRAP
UniProt Entry Name	CD40L_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/cd40lg-human-elisa-kits/scd40l/45308>