

Rabbit anti-Mouse TARC Polyclonal Antibody | anti-CCL17 antibody

Anti-TARC Antibody

CATALOG # AAA45702	HOST Rabbit	REACTIVITY Mouse	APPLICATIONS ELISA, Western Blot
------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

TARC; Polyclonal Antibody; Anti-TARC Antibody; C-C motif chemokine 17; chemokine (C-C motif) ligand 17; A-152E5.3 antibody; A152E53 antibody; ABCD 2 antibody; ABCD2 antibody; C-C motif chemokine 17 antibody; CC chemokine TARC antibody; CCL17 antibody; CCL17_HUMAN antibody; Chemokine CC Motif Ligand 17 antibody; MGC138271 antibody; MGC138273 antibody; SCYA17 antibody; Small Inducible Cytokine A17 antibody; Small Inducible Cytokine A17 Precursor antibody; Small Inducible Cytokine Subfamily A (Cys Cys) antibody; Small Inducible Cytokine Subfamily A (Cys Cys) Member 17 antibody; Small-inducible cytokine A17 antibody; T Cell Directed CC Chemokine antibody; Thymus and activation regulated chemokine antibody; Thymus and activation-regulated chemokine antibody; anti-CCL17 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	No cross reactivity with other proteins.
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized
Sequence Length	93
Applicable Applications for anti-CCL17 antibody	ELISA, WB (Western Blot)
Immunogen	E Coli-derived mouse TARC recombinant protein (Position: A34-P103). Mouse TARC shares 71% amino acid (aa) sequence identity with human TARC.
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Reconstitution	0.2ml of distilled water will yield a concentration of 500ug/ml.
Preparation and Storage	At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

IMAGES



Application Data
Anti-TARC Picoband antibody, AAA45702-1.jpgAll lanes: Anti-TARC(AAA45702) at 0.5ug/mlWB: Recombinant Mouse TARC Protein 0.5ngPredicted bind size: 34KDObserved bind size: 34KD

ADDITIONAL INFORMATION

Related Product Information for anti-CCL17 antibody	Description: Rabbit IgG polyclonal antibody for C-C motif chemokine 17(CCL17) detection. Tested with WB, ELISA in Mouse. Background: Chemokine (C-C motif) ligand 17 (CCL17) is a small cytokine belonging to the CC chemokine family that is also known as thymus and activation regulated chemokine (TARC). CCL17 is expressed constitutively in thymus, but only transiently in phytohemagglutinin-stimulated peripheral blood mononuclear cells. This chemokine specifically binds and induces chemotaxis in T cells and elicits its effects by interacting with the chemokine receptor CCR4. The gene for CCL17 is located on chromosome 16, in humans, along with other chemokines called CCL22 and CX3CL1. The standard used in this kit is recombinant human CCL17, consisting of 71 amino acids with the molecular weight of 8Kda.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	81907638
NCBI GeneID	20295
UniProt Accession #	Q9WUZ6
Molecular Weight	10,467 Da
NCBI Official Full Name	Thymus and activation-regulated chemokine (Chemokine (C-C motif) ligand 17) (10 day old male pancreas cDNA, RIKEN full-length enriched library, clone:1810031C02 product:small inducible cytokine subfamily A17, full insert sequence) (CC chemokine ABCD-2)
NCBI Official Synonym Full Names	chemokine (C-C motif) ligand 17
NCBI Official Symbol	Ccl17
NCBI Official Synonym Symbols	Tarc; Abcd-2; Scya17; Scya17l
NCBI Protein Information	chemokine (C-C motif) ligand 17; chemokine (C-C motif) ligand 17; small inducible cytokine subfamily A17
UniProt Protein Name	C-C motif chemokine
UniProt Gene Name	Ccl17
UniProt Entry Name	Q9WUZ6_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.