

Human ALCAM ELISA Kit | ALCAM elisa kit

Human ALCAM PicoKine ELISA Kit

CATALOG #
AAA45912

REACTIVITY
Human


PRODUCT SYNONYMS

ALCAM; Human ALCAM PicoKine ELISA Kit; CD166 antigen; activated leukocyte cell adhesion molecule; Activated leukocyte cell adhesion molecule; ALCAM protein; CD 166; CD166; CD166_HUMAN; FLJ3851; FLJ38514; MEMD; MGC71733; ALCAM elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	Natural and recombinant human ALCAM
Sequence Length	583
Samples	Cell culture supernates, serum and plasma (heparin, EDTA)
Assay Type	Sandwich
Detection Range	312pg/ml-20,000pg/ml
Sensitivity	< 10pg/ml
Intra-assay Precision	Intra-Assay Precision (Precision within an assay) Three samples of known concentration were tested on one plate to assess intra-assay precision.
Inter-assay Precision	Inter-Assay Precision (Precision between assays) Three samples of known concentration were tested in separate assays to assess inter-assay precision.
Preparation and Storage	Store at 4 degree C for 6 months, at -20 degree C for 12 months. Avoid multiple freeze-thaw cycles (Shipped with wet ice.)

IMAGES



Standard Curve (Sample)

Human ALCAM PicoKine ELISA Kit standard curve

ADDITIONAL INFORMATION

Related Product Information for ALCAM elisa kit	Principle of the assay: human ALCAM ELISA Kit was based on standard sandwich enzyme-linked immune-sorbent assay technology. A monoclonal antibody from mouse specific for ALCAM has been precoated onto 96-well plates. Standards(NSO, W28-A526) and test samples are added to the wells; a biotinylated detection polyclonal antibody from goat specific for ALCAM is added subsequently and then followed by washing with PBS or TBS buffer. Avidin-Biotin-Peroxidase Complex was added and unbound conjugates were washed away with PBS or TBS buffer. HRP substrate TMB was used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the human ALCAM amount of sample captured in plate. Background: CD166 antigen is a 100-105 kD typeI transmembrane glycoprotein that is a member of the immunoglobulin superfamily of proteins. In humans it is encoded by the ALCAM gene. The gene was mapped to human chromosome 3q13.1-q13.2 by fluorescence in situ hybridization of cDNA probes to metaphase chromosomes. It is also called CD166 (cluster of differentiation 166), MEMD, SC-1/DM-GRASP/BEN in the chicken, and KG-CAM in the rat. It is expressed on activated T cells, activated monocytes, epithelial cells, fibroblasts, neurons, melanoma cells, and also in sweat and sebaceous glands. CD166 protein expression is reported to be upregulated in a cell line deriving from a metastasizing melanoma. CD166 plays an important role in mediating adhesion interactions between thymic epithelial cells and CD6+ cells during intrathymic T cell development. Recently, CD166 has also been used as a potential cancer stem cell marker.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	118572629
NCBI GeneID	214
UniProt Accession #	Q13740; O60892; Q1HGM8; Q1HGM9; Q6PEY4; Q6ZS95; B2RNS3; B4DTU0
Molecular Weight	33,878 Da
NCBI Official Full Name	CD166 antigen
NCBI Official Synonym Full Names	activated leukocyte cell adhesion molecule
NCBI Official Symbol	ALCAM
NCBI Official Synonym Symbols	MEMD; CD166
NCBI Protein Information	CD166 antigen
UniProt Protein Name	CD166 antigen
UniProt Gene Name	ALCAM
UniProt Synonym Gene Names	MEMD
UniProt Entry Name	CD166_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/alcam/45912>