

Rabbit ERK1 Monoclonal Antibody | anti-MAPK3 antibody

Anti-ERK1 Rabbit Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA46873	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry

PRODUCT SYNONYMS

ERK1; Monoclonal Antibody; Anti-ERK1 Rabbit Monoclonal Antibody; Mitogen-activated protein kinase 3; MAP kinase 3; MAPK 3; 2.7.11.24; ERT2; Extracellular signal-regulated kinase 1; ERK-1; Insulin-stimulated MAP2 kinase; MAP kinase isoform p44; p44-MAPK; Microtubule-associated protein 2 kinase; p44-ERK1; MAPK3; PRKM3; anti-MAPK3 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	357
Applicable Applications for anti-MAPK3 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Immunogen	A synthesized peptide derived from human ERK1
Subcellular Localization	Cytoplasm. Nucleus. Autophosphorylation at Thr-207 promotes nuclear localization.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



WB (Western Blot)

Western blot analysis of ERK1 in Jurkat cell lysate (AAA46873). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MAPK3 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for MAPK3



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human bladder, using ERK1 Antibody(AAA46873)MAPK3 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-MAPK3 Antibody (AAA46873)overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-MAPK3 antibody	Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. MAPK1/ERK2 and MAPK3/ERK1 are the 2 MAPKs which play an important role in the MAPK/ERK cascade. They participate also in a signaling cascade initiated by activated KIT and KITLG/SCF. Depending on the cellular context, the MAPK/ERK cascade mediates diverse biological functions such as cell growth, adhesion, survival and differentiation through the regulation of transcription, translation, cytoskeletal rearrangements. The MAPK/ERK cascade plays also a role in initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors. About 160 substrates have already been discovered for ERKs. Many of these substrates are localized in the nucleus, and seem to participate in the regulation of transcription upon stimulation. However, other substrates are found in the cytosol as well as in other cellular organelles, and those are responsible for processes such as translation, mitosis and apoptosis. Moreover, the MAPK/ERK cascade is also involved in the regulation of the endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC); as well as in the fragmentation of the Golgi apparatus during mitosis. The substrates include transcription factors (such as ATF2, BCL6, ELK1, ERF, FOS, HSF4 or SPZ1), cytoskeletal elements (such as CANX, CTTN, GJA1, MAP2, MAPT, PXN, SORBS3 or STMN1), regulators of apoptosis (such as BAD, BTG2, CASP9, DAPK1, IER3, MCL1 or PPARG), regulators of translation (such as EIF4EBP1) and a variety of other signaling-related molecules (like ARHGEF2, FRS2 or GRB10). Protein kinases (such as RAF1, RPS6KA1/RSK1, RPS6KA3/RSK2, RPS6KA2/RSK3, RPS6KA6/RSK4, SYK, MKNK1/MNK1, MKNK2/MNK2, RPS6KA5/MSK1, RPS6KA4/MSK2, MAPKAPK3 or MAPKAPK5) and phosphatases (such as DUSP1, DUSP4, DUSP6 or DUSP16) are other substrates which enable the propagation the MAPK/ERK signal to additional cytosolic and nuclear targets, thereby extending the specificity of the cascade.
Product Categories/Family for anti-MAPK3 antibody	Neuroscience; Neurology Process; Neurodegenerative Disease; Alzheimer's Disease; tangles & tau; Signal Transduction; Protein Phosphorylation; SER/THR Kinases; MAPK Pathway; Stem Cells; Signaling Pathways; TGF Beta; Cytoplasmic; Kits/Lysates/Other; ELISA K

NCBI AND UNIPROT INFORMATION

NCBI GI #	91718897
NCBI GeneID	5595
NCBI Accession #	NP_001035145.1
NCBI GenBank Nucleotide #	NM_001040056.2
UniProt Accession #	P27361; Q8NHX1; A8CZ58; B0LPG3
Molecular Weight	43136 MW
NCBI Official Full Name	mitogen-activated protein kinase 3 isoform 2
NCBI Official Synonym Full Names	mitogen-activated protein kinase 3
NCBI Official Symbol	MAPK3
NCBI Official Synonym Symbols	ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK
NCBI Protein Information	mitogen-activated protein kinase 3
UniProt Protein Name	Mitogen-activated protein kinase 3
UniProt Gene Name	MAPK3
UniProt Synonym Gene Names	ERK1; PRKM3; MAP kinase 3; MAPK 3; ERK-1; p44-MAPK

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/erk1-antibody-western-blot-immunohistochemistry-elisa/46873>