Rabbit anti-Human Bcl-6 Monoclonal Antibody | anti-Bcl-6 antibody

Anti-Bcl-6 Rabbit Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA46884	**Rabbit**	**Human**	**Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence, Immunoprecipitation**

PRODUCT SYNONYMS

Bcl-6; Monoclonal Antibody; Anti-Bcl-6 Rabbit Monoclonal Antibody; B-cell lymphoma 6 protein; BCL-6; B-cell lymphoma 5 protein; BCL-5; Protein LAZ-3; Zinc finger and BTB domain-containing protein 27; Zinc finger protein 51; BCL6; BCL5; LAZ3; ZBTB27; ZNF51; anti-Bcl-6 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	650
Applicable Applications for anti-Bcl-6 antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence), IP (Immunoprecipitation)
Immunogen	A synthesized peptide derived from human Bcl6
Subcellular Localization	Nucleus.
Tissue Specificity	Expressed in germinal center T- and B-cells and in primary immature dendritic cells.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES

WB (Western Blot)
Western blot analysis of Bcl6 in expression Daudi cell lysate (AAA46884).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-BCL6 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for BCL6

IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human tonsil, using Bcl-6 Antibody(AAA46884)BCL6 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-BCL6 Antibody (AAA46884)overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-Bcl-6 antibody	Transcriptional repressor mainly required for germinal center (GC) formation and antibody affinity maturation which has different mechanisms of action specific to the lineage and biological functions. Forms complexes with different corepressors and histone deacetylases to repress the transcriptional expression of different subsets of target genes. Represses its target genes by binding directly to the DNA sequence 5'-TTCCTAGAA-3' (BCL6- binding site) or indirectly by repressing the transcriptional activity of transcription factors. In GC B-cells, represses genes that function in differentiation, inflammation, apoptosis and cell cycle control, also autoregulates its transcriptional expression and up-regulates, indirectly, the expression of some genes important for GC reactions, such as AICDA, through the repression of microRNAs expression, like miR155. An important function is to allow GC B-cells to proliferate very rapidly in response to T-cell dependent antigens and tolerate the physiological DNA breaks required for immunglobulin class switch Recombinantion and somatic hypermutation without inducing a p53/TP53-dependent apoptotic response. In follicular helper CD4(+) T-cells (T(FH) cells), promotes the expression of T(FH)-related genes but inhibits the differentiation of T(H)1, T(H)2 and T(H)17 cells. Also required for the establishment and maintenance of immunological memory for both T- and B-cells. Suppresses macrophage proliferation through competition with STAT5 for STAT-binding motifs binding on certain target genes, such as CCL2 and CCND2. In response to genotoxic stress, controls cell cycle arrest in GC B-cells in both p53/TP53- dependedent and -independent manners. Besides, also controls neurogenesis through the alteration of the composition of NOTCH- dependent transcriptional complexes at selective NOTCH targets, such as HES5, including the recruitment of the deacetylase SIRT1 and resulting in an epigenetic silencing leading to neuronal differentiation.
Product Categories/Family for anti-Bcl-6 antibody	Epigenetics and Nuclear Signaling; Transcription; Domain Families; zinc finger; Epigenetics and Nuclear Signaling; Cancer Susceptibility; Proto-Oncogenes; Cancer; Oncoproteins/Suppressors; Oncoproteins; Transcription Factors

NCBI AND UNIPROT INFORMATION

NCBI GI #	195927002
NCBI GeneID	604
NCBI Accession #	NP_001124317.1
NCBI GenBank Nucleotide #	NM_001130845.1
UniProt Accession #	P41182; A7E241; B8PSA7; D3DNV5
Molecular Weight	78846 MW
NCBI Official Full Name	B-cell lymphoma 6 protein isoform 1
NCBI Official Synonym Full Names	B cell CLL/lymphoma 6
NCBI Official Symbol	BCL6
NCBI Official Synonym Symbols	BCL5; LAZ3; BCL6A; ZNF51; ZBTB27
NCBI Protein Information	B-cell lymphoma 6 protein
UniProt Protein Name	B-cell lymphoma 6 protein
UniProt Gene Name	**BCL6**
UniProt Synonym Gene Names	BCL5; LAZ3; ZBTB27; ZNF51; BCL-6; BCL-5

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/monoclonal/bcl-6-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-immunoprecipitation-elisa/46884