

Rabbit Cyclin D1 Monoclonal Antibody | anti-CCND1 antibody

Anti-Cyclin D1 Rabbit Monoclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA46888	Rabbit	Human, Mouse, Rat	Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence, Immunoprecipitation

PRODUCT SYNONYMS


Cyclin D1; Monoclonal Antibody; Anti-Cyclin D1 Rabbit Monoclonal Antibody; G1/S-specific cyclin-D1; B-cell lymphoma 1 protein; BCL-1; BCL-1 oncogene; PRAD1 oncogene; CCND1; BCL1; PRAD1; anti-CCND1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	295
Applicable Applications for anti-CCND1 antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence), IP (Immunoprecipitation)
Immunogen	A synthesized peptide derived from human Cyclin D1
Subcellular Localization	Nucleus. Cytoplasm. Membrane. Cyclin D-CDK4 complexes accumulate at the nuclear membrane and are then translocated to the nucleus through interaction with KIP/CIP family members.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



IHC (Immunohiostchemistry)
Immunohistochemical analysis of paraffin-embedded human bladder, using Cyclin D1 Antibody(AAA46888)CCND1 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CCND1 Antibody (AAA46888)overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)
Western blot analysis of Cyclin D1 expression in (1)MCF-7 cell lysates;(2) LnCaP cell lysates (AAA46888).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CCND1 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for CCND1

ADDITIONAL INFORMATION

Related Product Information for anti-CCND1 antibody	Regulatory component of the cyclin D1-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals. Also substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and repressing its transcriptional activity. Component of the ternary complex, cyclin D1/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex. Exhibits transcriptional corepressor activity with INSM1 on the NEUROD1 and INS promoters in a cell cycle-independent manner.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	16950655
NCBI GeneID	595
NCBI Accession #	NP_444284.1
NCBI GenBank Nucleotide #	NM_053056.2
UniProt Accession #	P24385; Q6LEF0
Molecular Weight	33729 MW
NCBI Official Full Name	G1/S-specific cyclin-D1
NCBI Official Synonym Full Names	cyclin D1
NCBI Official Symbol	CCND1
NCBI Official Synonym Symbols	BCL1; PRAD1; U21B31; D11S287E
NCBI Protein Information	G1/S-specific cyclin-D1
UniProt Protein Name	G1/S-specific cyclin-D1

UniProt Gene Name	CCND1
UniProt Synonym Gene Names	BCL1; PRAD1; BCL-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cyclin-d1-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-immunoprecipitation-elisa/46888>