

Rabbit PI 3 Kinase p85 alpha Monoclonal Antibody | anti-PIK3R1 antibody

Anti-PI 3 Kinase p85 alpha Rabbit Monoclonal Antibody

CATALOG # AAA46926	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

PI 3 Kinase p85 alpha; Monoclonal Antibody; Anti-PI 3 Kinase p85 alpha Rabbit Monoclonal Antibody; Phosphatidylinositol 3-kinase regulatory subunit alpha; PI3-kinase regulatory subunit alpha; PI3K regulatory subunit alpha; PtdIns-3-kinase regulatory subunit alpha; Phosphatidylinositol 3-kinase 85 kDa regulatory subunit alpha; PI3-kinase subunit p85-alpha; PtdIns-3-kinase regulatory subunit p85-alpha; PIK3R1; GRB1; anti-PIK3R1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Concentration	1 mg/mL (varies by lot)
Sequence Length	361
Applicable Applications for anti-PIK3R1 antibody	WB (Western Blot)
Immunogen	A synthesized peptide derived from human PI 3 Kinase p85 alpha
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



WB (Western Blot)
Western blot analysis of PI 3 Kinase p85 alpha expression in A431 cell lysate (AAA46926). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PIK3R1 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for PIK3R1.

ADDITIONAL INFORMATION

Related Product Information for anti-PIK3R1 antibody	Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues. Plays an important role in signaling in response to FGFR1, FGFR2, FGFR3, FGFR4, KITLG/SCF, KIT, PDGFRA and PDGFRB. Likewise, plays a role in ITGB2 signaling (PubMed: 17626883, PubMed: 19805105, PubMed: 7518429). Modulates the cellular response to ER stress by promoting nuclear translocation of XBP1 isoform 2 in a ER stress- and/or insulin-dependent manner during metabolic overloading in the liver and hence plays a role in glucose tolerance improvement (PubMed: 20348923).
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	335057534
NCBI GeneID	5295
NCBI Accession #	NP_001229395.1
NCBI GenBank Nucleotide #	NM_001242466.1
UniProt Accession #	P27986; Q15747; Q4VBZ7; Q53EM6; Q8IXA2; Q8N1C5; B3KT19; D3DWA0; E7EX19
NCBI Official Full Name	phosphatidylinositol 3-kinase regulatory subunit alpha isoform 4
NCBI Official Synonym Full Names	phosphoinositide-3-kinase regulatory subunit 1
NCBI Official Symbol	PIK3R1
NCBI Official Synonym Symbols	p85; AGM7; GRB1; IMD36; p85-ALPHA
NCBI Protein Information	phosphatidylinositol 3-kinase regulatory subunit alpha
UniProt Protein Name	Phosphatidylinositol 3-kinase regulatory subunit alpha
UniProt Gene Name	PIK3R1
UniProt Synonym Gene Names	GRB1; PI3-kinase regulatory subunit alpha; PI3K regulatory subunit alpha; PtdIns-3-kinase regulatory subunit alpha; PI3-kinase subunit p85-alpha; PtdIns-3-kinase regulatory subunit p85-alpha

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/pi-3-kinase-p85-alpha-antibody-western-blot-elisa/46926>