

Rabbit anti-Human CD46 Monoclonal Antibody | anti-CD46 antibody

Anti-CD46 Rabbit Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA46937	Rabbit	Human	Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence

PRODUCT SYNONYMS

CD46; Monoclonal Antibody; Anti-CD46 Rabbit Monoclonal Antibody; Membrane cofactor protein; TLX; Trophoblast leukocyte common antigen; MCP; MIC10; anti-CD46 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	336
Applicable Applications for anti-CD46 antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence)
Immunogen	A synthesized peptide derived from human CD46
Subcellular Localization	Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane; Single- pass type I membrane protein. Inner acrosomal membrane of spermatozoa. Internalized upon binding of Measles virus, Herpesvirus 6 or Neisseria gonorrhoeae, which results in an increased susceptibility of infected cells to complement-mediated injury. In cancer cells or cells infected by Neisseria, shedding leads to a soluble peptide.
Tissue Specificity	Expressed by all cells except erythrocytes.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



WB (Western Blot)

Western blot analysis of CD46 in K562 cell lysate (AAA46937). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CD46 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for CD46



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human colon, using CD46 Antibody(AAA46937). CD46 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CD46 Antibody (AAA46937) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-CD46 antibody	Acts as a cofactor for complement factor I, a serine protease which protects autologous cells against complement-mediated injury by cleaving C3b and C4b deposited on host tissue. May be involved in the fusion of the spermatozoa with the oocyte during fertilization. Also acts as a costimulatory factor for T- cells which induces the differentiation of CD4+ into T-regulatory 1 cells. T-regulatory 1 cells suppress immune responses by secreting interleukin-10, and therefore are thought to prevent autoimmunity. A number of viral and bacterial pathogens seem to exploit this property and directly induce an immunosuppressive phenotype in T-cells by binding to CD46.
Product Categories/Family for anti-CD46 antibody	Immunology; Cell Type Markers; Non-Lineage; Immunology; Innate Immunity; complement; regulatory

NCBI AND UNIPROT INFORMATION

NCBI GI #	27502402
NCBI GeneID	4179
NCBI Accession #	NP_002380.3
NCBI GenBank Nucleotide #	NM_002389.4
UniProt Accession #	P15529; Q15429; Q53GV9; Q5HY94; Q5VWS6; Q5VWS7; Q5VWS8; Q5VWS9; Q5VWT0; A0T1T0; A0T1T1; A0T1T2
Molecular Weight	43747 MW
NCBI Official Full Name	membrane cofactor protein isoform 1
NCBI Official Synonym Full Names	CD46 molecule
NCBI Official Symbol	CD46
NCBI Official Synonym Symbols	MCP; TLX; AHUS2; MIC10; TRA2.10
NCBI Protein Information	membrane cofactor protein
UniProt Protein Name	Membrane cofactor protein
UniProt Gene Name	CD46
UniProt Synonym Gene Names	MCP; MIC10

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd46-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-elisa/46937>