

Rabbit Integrin beta 3 Monoclonal Antibody | anti-ITGB3 antibody

Anti-Integrin beta 3 Rabbit Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA46962	Rabbit	Human, Mouse, Rat	Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence


PRODUCT SYNONYMS

Integrin beta 3; Monoclonal Antibody; Anti-Integrin beta 3 Rabbit Monoclonal Antibody; Integrin beta-3; Platelet membrane glycoprotein IIIa; GPIIIa; CD61; ITGB3; GP3A; anti-ITGB3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	784
Applicable Applications for anti-ITGB3 antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence)
Immunogen	A synthesized peptide derived from human Integrin beta 3
Subcellular Localization	Cell membrane; Single- pass type I membrane protein. Cell projection, lamellipodium membrane. Cell junction, focal adhesion.
Tissue Specificity	Isoform beta-3A and isoform beta-3C are widely expressed. Isoform beta-3A is specifically expressed in osteoblast cells; isoform beta-3C is specifically expressed in prostate and testis.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human testis, using Integrin beta 3 Antibody (AAA46962) ITGB3 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-ITGB3 Antibody (AAA46962) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Western blot analysis of Integrin beta3 expression in human spleen lysate (AAA46962). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ITGB3 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ITGB3

ADDITIONAL INFORMATION

Related Product Information for anti-ITGB3 antibody	Integrin alpha-V/beta-3 (ITGAV: ITGB3) is a receptor for cytotactin, fibronectin, laminin, matrix metalloproteinase-2, osteopontin, osteomodulin, prothrombin, thrombospondin, vitronectin and von Willebrand factor. Integrin alpha-IIb/beta-3 (ITGA2B: ITGB3) is a receptor for fibronectin, fibrinogen, plasminogen, prothrombin, thrombospondin and vitronectin. Integrins alpha-IIb/beta-3 and alpha-V/beta-3 recognize the sequence R-G-D in a wide array of ligands. Integrin alpha- IIb/beta-3 recognizes the sequence H-H-L-G-G-G-A-K-Q-A-G-D-V in fibrinogen gamma chain. Following activation integrin alpha- IIb/beta-3 brings about platelet/platelet interaction through binding of soluble fibrinogen. This step leads to rapid platelet aggregation which physically plugs ruptured endothelial surface. Fibrinogen binding enhances SELP expression in activated platelets (By similarity). In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions.
Product Categories/Family for anti-ITGB3 antibody	Epigenetics and Nuclear Signaling; Chromosome Structure; centromeres; Signal Transduction; Cytoskeleton/ECM; Cell Adhesion; Integrins; Signaling Pathway; NFKB Pathway; nuclear hormone Receptors; Co-Activators/Co-Repressors; Epigenetics and Nuclear Signali

NCBI AND UNIPROT INFORMATION

NCBI GI #	47078292
NCBI GeneID	3690
NCBI Accession #	NP_000203.2
NCBI GenBank Nucleotide #	NM_000212.2
UniProt Accession #	P05106; O15495; Q12806; Q13413; Q14648; Q16499; A0PJW2; D3DXJ8
Molecular Weight	87058 MW
NCBI Official Full Name	integrin beta-3
NCBI Official Synonym Full Names	integrin subunit beta 3
NCBI Official Symbol	ITGB3
NCBI Official Synonym Symbols	GT; CD61; GP3A; BDPLT2; GPIIIa; BDPLT16
NCBI Protein Information	integrin beta-3
UniProt Protein Name	Integrin beta-3
UniProt Gene Name	ITGB3
UniProt Synonym Gene Names	GP3A; GPIIIa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/integrin-beta-3-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-elisa/46962>