

Rabbit anti-Human CDX2 Monoclonal Antibody | anti-CDX2 antibody

Anti-CDX2 Rabbit Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA46998	Rabbit	Human	Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

CDX2; Monoclonal Antibody; Anti-CDX2 Rabbit Monoclonal Antibody; Homeobox protein CDX-2; CDX-3; Caudal-type homeobox protein 2; CDX3; anti-CDX2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	313
Applicable Applications for anti-CDX2 antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Immunogen	A synthesized peptide derived from human CDX2
Subcellular Localization	Nucleus.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



WB (Western Blot)

Western blot analysis of CDX2 in (1) HEK293 cell lysate; (2) Colon cancer lysate (AAA46998). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CDX2 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for CDX2



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human colon, using CDX2 Antibody(AAA46998). CDX2 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CDX2 Antibody (AAA46998) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-CDX2 antibody	Involved in the transcriptional regulation of multiple genes expressed in the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestine.
Product Categories/Family for anti-CDX2 antibody	Stem Cells; Lineage Markers; Endoderm; Epigenetics and Nuclear Signaling; Transcription; Transcription Factors; Developmental Biology; Lineage Specification; Trophectoderm

NCBI AND UNIPROT INFORMATION

NCBI GI #	372624392
NCBI GeneID	1045
NCBI Accession #	NP_001256.3
NCBI GenBank Nucleotide #	NM_001265.5
UniProt Accession #	Q99626; O00503; Q5VTU7; Q969L8; Q9UD92
Molecular Weight	33520 MW
NCBI Official Full Name	homeobox protein CDX-2 isoform 1
NCBI Official Synonym Full Names	caudal type homeobox 2
NCBI Official Symbol	CDX2
NCBI Official Synonym Symbols	CDX3; CDX-3; CDX2/AS
NCBI Protein Information	homeobox protein CDX-2
UniProt Protein Name	Homeobox protein CDX-2
UniProt Gene Name	CDX2
UniProt Synonym Gene Names	CDX3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.