

Rabbit GRP78 BiP Monoclonal Antibody | anti-GRP78 BiP antibody

Anti-GRP78 BiP Rabbit Monoclonal Antibody

CATALOG # AAA47009	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	--	---

PRODUCT SYNONYMS

GRP78 BiP; Monoclonal Antibody; Anti-GRP78 BiP Rabbit Monoclonal Antibody; 78 kDa glucose-regulated protein; GRP-78; Endoplasmic reticulum lumenal Ca(2+)-binding protein grp78; Heat shock 70 kDa protein 5; Immunoglobulin heavy chain-binding protein; BiP; HSPA5; GRP78; anti-GRP78 BiP antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	654
Applicable Applications for anti-GRP78 BiP antibody	WB (Western Blot), IHC (Immunohistochemistry)
Immunogen	A synthesized peptide derived from human GRP78 BiP
Subcellular Localization	Endoplasmic reticulum lumen. Melanosome. Cytoplasm. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



WB (Western Blot)

Western blot analysis of GRP78 BiP expression in (1) HeLa cell lysate; (2) C6 cell lysate (AAA47009). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-HSPA5 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for HSPA5



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human testis, using GRP78 BiP Antibody (AAA47009). HSPA5 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-HSPA5 Antibody (AAA47009) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-GRP78 BiP antibody	Probably plays a role in facilitating the assembly of multimeric protein complexes inside the endoplasmic reticulum. Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10, probably to facilitate the release of DNAJC10 from its substrate.
Product Categories/Family for anti-GRP78 BiP antibody	Tags & Cell Markers; Subcellular Markers; organelles

NCBI AND UNIPROT INFORMATION

NCBI GI #	16507237
NCBI GeneID	3309
NCBI Accession #	NP_005338.1
NCBI GenBank Nucleotide #	NM_005347.4
UniProt Accession #	P11021; Q2EF78; Q9NPF1; Q9UK02; B0QZ61
Molecular Weight	72333 MW
NCBI Official Full Name	78 kDa glucose-regulated protein
NCBI Official Synonym Full Names	heat shock protein family A (Hsp70) member 5
NCBI Official Symbol	HSPA5
NCBI Official Synonym Symbols	BIP; MIF2; GRP78; HEL-S-89n
NCBI Protein Information	78 kDa glucose-regulated protein
UniProt Protein Name	78 kDa glucose-regulated protein
UniProt Gene Name	HSPA5
UniProt Synonym Gene Names	BiP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.