

Rabbit NGFR/P75 Monoclonal Antibody | anti-NGFR antibody

Anti-NGFR/P75 Rabbit Monoclonal Antibody



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA47033	Rabbit	Human, Mouse, Rat	Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence, Immunoprecipitation

PRODUCT SYNONYMS

NGFR/P75; Monoclonal Antibody; Anti-NGFR/P75 Rabbit Monoclonal Antibody; Tumor necrosis factor receptor superfamily member 16; Gp80-LNGFR; Low affinity neurotrophin receptor p75NTR; Low-affinity nerve growth factor receptor; NGF receptor; p75 ICD; CD271; NGFR; TNFRSF16; anti-NGFR antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	333
Applicable Applications for anti-NGFR antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence), IP (Immunoprecipitation)
Immunogen	A synthesized peptide derived from human p75 NGF Receptor
Subcellular Localization	Membrane; Single-pass type I membrane protein.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human glioma, using NGFR Antibody (AAA47033)NGFR was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-NGFR Antibody (AAA47033)overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

WB (Western Blot)
Western blot analysis of NGFR expression in (1) C6 cell lysate; (2) PC-12 cell lysate (AAA47033).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-NGFR monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for NGFR

ADDITIONAL INFORMATION

Related Product Information for anti-NGFR antibody	Plays a role in the regulation of the translocation of GLUT4 to the cell surface in adipocytes and skeletal muscle cells in response to insulin, probably by regulating RAB31 activity, and thereby contributes to the regulation of insulin-dependent glucose uptake (By similarity). Low affinity receptor which can bind to NGF, BDNF, NT-3, and NT-4. Can mediate cell survival as well as cell death of neural cells. Necessary for the circadian oscillation of the clock genes ARNTL/BMAL1, PER1, PER2 and NR1D1 in the suprachiasmatic nucleus (SCN) of the brain and in liver and of the genes involved in glucose and lipid metabolism in the liver.
Product Categories/Family for anti-NGFR antibody	Neuroscience; Neurology Process; Growth and Development; neurotrophins; Stem Cells; Neural Stem Cells; Surface Molecules; mesenchymal stem cells

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505393
NCBI GeneID	4804
NCBI Accession #	NP_002498.1
NCBI GenBank Nucleotide #	NM_002507.3
UniProt Accession #	P08138; B2R961; B4E096
Molecular Weight	45183 MW
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 16
NCBI Official Synonym Full Names	nerve growth factor receptor
NCBI Official Symbol	NGFR
NCBI Official Synonym Symbols	CD271; p75NTR; TNFRSF16; p75(NTR); Gp80-LNGFR
NCBI Protein Information	tumor necrosis factor receptor superfamily member 16
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 16
UniProt Gene Name	NGFR

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/ngfr-p75-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-immunoprecipitation-elisa/47033>