

Rabbit Stathmin 1 Monoclonal Antibody | anti-STMN1 antibody

Anti-Stathmin 1 Rabbit Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA47035	Rabbit	Human, Mouse, Rat	Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence, Immunoprecipitation, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Stathmin 1; Monoclonal Antibody; Anti-Stathmin 1 Rabbit Monoclonal Antibody; Stathmin; Leukemia-associated phosphoprotein p18; Metablastin; Oncoprotein 18; Op18; Phosphoprotein p19; pp19; Prosolin; Protein Pr22; pp17; STMN1; C1orf215; LAP18; OP18; anti-STMN1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	174
Applicable Applications for anti-STMN1 antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence), IP (Immunoprecipitation), FCM/FACS (Flow Cytometry)
Immunogen	A synthesized peptide derived from human Stathmin 1
Subcellular Localization	Cytoplasm, cytoskeleton.
Tissue Specificity	Ubiquitous. Expression is strongest in fetal and adult brain, spinal cord, and cerebellum, followed by thymus, bone marrow, testis, and fetal liver. Expression is intermediate in colon, ovary, placenta, uterus, and trachea, and is readily detected at substantially lower levels in all other tissues examined. Lowest expression is found in adult liver. Present in much greater abundance in cells from patients with acute leukemia of different subtypes than in normal peripheral blood lymphocytes, non-leukemic proliferating lymphoid cells, bone marrow cells, or cells from patients with chronic lymphoid or myeloid leukemia.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human colon, using Stathmin 1 Antibody(AAA47035)STMN1 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-STMN1 Antibody (AAA47035)overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)
Western blot analysis of Stathmin 1 expression in PC-12 cell lysate (AAA47035).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-STMN1 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for STMN1

ADDITIONAL INFORMATION

Related Product Information for anti-STMN1 antibody	Involved in the regulation of the microtubule (MT) filament system by destabilizing microtubules. Prevents assembly and promotes disassembly of microtubules. Phosphorylation at Ser- 16 may be required for axon formation during neurogenesis. Involved in the control of the learned and innate fear (By similarity).
Product Categories/Family for anti-STMN1 antibody	Neuroscience; Neurology Process; Neurodegenerative Disease; Alzheimer's Disease; Neuroscience; cell adhesion proteins; cytoskeletal proteins; microtubules; Signal Transduction; Cytoskeleton/ECM; Cytoskeleton; tubulin

NCBI AND UNIPROT INFORMATION

NCBI GI #	224451142
NCBI GeneID	3925
NCBI Accession #	NP_001138926.1
NCBI GenBank Nucleotide #	NM_001145454.2
UniProt Accession #	P16949; A2A2D1; B2R4E7; B7Z8N4; D3DPJ5
Molecular Weight	17303 MW
NCBI Official Full Name	stathmin isoform b
NCBI Official Synonym Full Names	stathmin 1
NCBI Official Symbol	STMN1
NCBI Official Synonym Symbols	Lag; SMN; OP18; PP17; PP19; PR22; LAP18; C1orf215
NCBI Protein Information	stathmin
UniProt Protein Name	Stathmin
UniProt Gene Name	STMN1
UniProt Synonym Gene Names	C1orf215; LAP18; OP18; Op18; pp19

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/stathmin-1-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-immunoprecipitation-flow-cytometry-elisa/47035>