

Rabbit TGFB1/Beta Ig H3 Monoclonal Antibody | anti-TGFB1 antibody

Anti-TGFB1/Beta Ig H3 Rabbit Monoclonal Antibody

CATALOG # AAA47039	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	--	---

PRODUCT SYNONYMS

TGFB1/Beta Ig H3; Monoclonal Antibody; Anti-TGFB1/Beta Ig H3 Rabbit Monoclonal Antibody; Transforming growth factor-beta-induced protein ig-h3; Beta ig-h3; Kerato-epithelin; RGD-containing collagen-associated protein; RGD-CAP; TGFB1; BIGH3; anti-TGFB1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	683
Applicable Applications for anti-TGFB1 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Immunogen	A synthesized peptide derived from human TGFB1
Subcellular Localization	Secreted, extracellular space, extracellular matrix. May be associated both with microfibrils and with the cell surface.
Tissue Specificity	Highly expressed in the corneal epithelium.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



WB (Western Blot)

Western blot analysis of TGFB1 expression in human fetal kidney lysate (AAA47039). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TGFB1 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for TGFB1



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human kidney, using TGFB1 Antibody(AAA47039). TGFB1 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-TGFB1 Antibody (AAA47039) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-TGFBI antibody	Binds to type I, II, and IV collagens. This adhesion protein may play an important role in cell-collagen interactions. In cartilage, may be involved in endochondral bone formation.
Product Categories/Family for anti-TGFBI antibody	Signal Transduction; Cytoskeleton/ECM; Extracellular Matrix; ECM Proteins; Collagen; Signal Transduction; Growth Factors/Hormones; TGF; Structures; Bone; Neuroscience; Sensory System; Visual System; Other ECM Proteins; Developmental Biology; Organogenesis

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507467
NCBI GeneID	7045
NCBI Accession #	NP_000349.1
NCBI GenBank Nucleotide #	NM_000358.2
UniProt Accession #	Q15582; O14471; O14472; O14476; O43216; O43217; O43218; O43219; Q53XM1; D3DQB1
Molecular Weight	74681 MW
NCBI Official Full Name	transforming growth factor-beta-induced protein ig-h3
NCBI Official Synonym Full Names	transforming growth factor beta induced
NCBI Official Symbol	TGFBI
NCBI Official Synonym Symbols	CSD; CDB1; CDG2; CSD1; CSD2; CSD3; EBMD; LCD1; BIGH3; CDGG1
NCBI Protein Information	transforming growth factor-beta-induced protein ig-h3
UniProt Protein Name	Transforming growth factor-beta-induced protein ig-h3
UniProt Gene Name	TGFBI
UniProt Synonym Gene Names	BIGH3; Beta ig-h3; RGD-CAP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/tgfb-beta-ig-h3-antibody-western-blot-immunohistochemistry-elisa/47039>