

Rabbit anti-Human Caveolin-2 Monoclonal Antibody | anti-CAV2 antibody

Anti-Caveolin-2 Rabbit Monoclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA47069	Rabbit	Human	Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence, Immunoprecipitation

PRODUCT SYNONYMS


Caveolin-2; Monoclonal Antibody; Anti-Caveolin-2 Rabbit Monoclonal Antibody; CAV2; anti-CAV2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	112
Applicable Applications for anti-CAV2 antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence), IP (Immunoprecipitation)
Immunogen	A synthesized peptide derived from human Caveolin-2
Subcellular Localization	Nucleus. Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Cell membrane; Peripheral membrane protein. Membrane, caveola; Peripheral membrane protein. Potential hairpin-like structure in the membrane. Membrane protein of caveolae. Tyr-19-phosphorylated form is enriched at sites of cell-cell contact and is translocated to the nucleus in complex with MAPK1 in response to insulin (By similarity). Tyr-27- phosphorylated form is located both in the cytoplasm and plasma membrane. CAV1-mediated Ser-23-phosphorylated form locates to the plasma membrane. Ser-36-phosphorylated form resides in intracellular compartments.
Tissue Specificity	Expressed in endothelial cells, smooth muscle cells, skeletal myoblasts and fibroblasts.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human colon, using Caveolin-2 Antibody(AAA47069)CAV2 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CAV2 Antibody (AAA47069)overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)
Western blot analysis of Caveolin 2 expression in HeLa cell lysate (AAA47069).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CAV2 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for CAV2

ADDITIONAL INFORMATION

Related Product Information for anti-CAV2 antibody	May act as a scaffolding protein within caveolar membranes. Interacts directly with G-protein alpha subunits and can functionally regulate their activity. Acts as an accessory protein in conjunction with CAV1 in targeting to lipid rafts and driving caveolae formation. The Ser-36 phosphorylated form has a role in modulating mitosis in endothelial cells. Positive regulator of cellular mitogenesis of the MAPK signaling pathway. Required for the insulin-stimulated nuclear translocation and activation of MAPK1 and STAT3, and the subsequent regulation of cell cycle progression (By similarity).
Product Categories/Family for anti-CAV2 antibody	Neuroscience; Neurotransmission; Calcium Signaling; calcium channels; p/q/r -type; Signal Transduction; Signaling Pathway; Cancer; Cancer Metabolism; Metabolic Signaling Pathway; Integration of Energy Metabolism; Metabolism; Pathways and Processes; Metabo

NCBI AND UNIPROT INFORMATION

NCBI GI #	332164664
NCBI GeneID	858
NCBI Accession #	NP_001193676.1
NCBI GenBank Nucleotide #	NM_001206747.1
UniProt Accession #	P51636; Q9UGM7; A4D0U2
Molecular Weight	18291 MW
NCBI Official Full Name	caveolin-2 isoform b
NCBI Official Synonym Full Names	caveolin 2
NCBI Official Symbol	CAV2
NCBI Official Synonym Symbols	CAV
NCBI Protein Information	caveolin-2
UniProt Protein Name	Caveolin-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/caveolin-2-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-immunoprecipitation-elisa/47069>