

Rabbit anti-Human, Rat Cytokeratin 20 Monoclonal Antibody | anti-KRT20 antibody

Anti-Cytokeratin 20 Rabbit Monoclonal Antibody



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA47121	Rabbit	Human, Rat	Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence, Immunoprecipitation, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

Cytokeratin 20; Monoclonal Antibody; Anti-Cytokeratin 20 Rabbit Monoclonal Antibody; Keratin; type I cytoskeletal 20; Cytokeratin-20; CK-20; Keratin-20; K20; Protein IT; KRT20; anti-KRT20 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	424
Applicable Applications for anti-KRT20 antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence), IP (Immunoprecipitation), FCM/FACS (Flow Cytometry)
Immunogen	A synthesized peptide derived from human Cytokeratin 20
Subcellular Localization	Cytoplasm.
Tissue Specificity	Expressed predominantly in the intestinal epithelium. Expressed in luminal cells of colonic mucosa. Also expressed in the Merkel cells of keratinized oral mucosa; specifically at the tips of some rete ridges of the gingival mucosa, in the basal layer of the palatal mucosa and in the taste buds of lingual mucosa.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human colon, using Cytokeratin 20 Antibody(AAA47121).KRT20 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-KRT20 Antibody (AAA47121)overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

WB (Western Blot)
Western blot analysis of Cytokeratin 20 expression in SW480 cell lysate (AAA47121).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-KRT20 monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for KRT20

ADDITIONAL INFORMATION

Related Product Information for anti-KRT20 antibody	Plays a significant role in maintaining keratin filament organization in intestinal epithelia. When phosphorylated, plays a role in the secretion of mucin in the small intestine (By similarity).
Product Categories/Family for anti-KRT20 antibody	Signal Transduction; Cytoskeleton/ECM; Cytoskeleton; Intermediate Filaments; class i; cytokeratins; Cancer; Tumor Biomarkers

NCBI AND UNIPROT INFORMATION

NCBI GI #	27894337
NCBI GeneID	54474
NCBI Accession #	NP_061883.1
NCBI GenBank Nucleotide #	NM_019010.2
UniProt Accession #	P35900; B2R6W7
Molecular Weight	48487 MW
NCBI Official Full Name	keratin, type I cytoskeletal 20
NCBI Official Synonym Full Names	keratin 20
NCBI Official Symbol	KRT20
NCBI Official Synonym Symbols	K20; CD20; CK20; CK-20; KRT21
NCBI Protein Information	keratin, type I cytoskeletal 20
UniProt Protein Name	Keratin, type I cytoskeletal 20
UniProt Gene Name	KRT20
UniProt Synonym Gene Names	CK-20; K20

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/cytokeratin-20-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-immunoprecipitation-flow-cytometry-elisa/47121>