

Rabbit anti-Human CD38 Monoclonal Antibody | anti-CD38 antibody

Anti-CD38 antibody (DM28), Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA47179	Rabbit	Human	Flow Cytometry, Functional Assay, ELISA


PRODUCT SYNONYMS

CD38; Monoclonal Antibody; Anti-CD38 antibody (DM28); Rabbit mAb; T10; cADPr hydrolase 1; anti-CD38 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	DM28
Purity/Purification	Purified from cell culture supernatant by affinity chromatography
Form/Format	Liquid; 0.1% Procline 300; 50% Glycerol; PBS, pH7.4; 0.1% BSA
Applicable Applications for anti-CD38 antibody	FCM/FACS (Flow Cytometry), ELISA
Immunogen	Recombinant human CD38 (Val43-Ile300) produced by using human HEK293 cells
Reconstitution	Reconstitute with deionized water
Preservative	0.1% Procline 300
Constituents	50% Glycerol; PBS,pH 7.4; 0.1% BSA
Preparation and Storage	Store at -20 degree C for 12 months (Avoid repeated freezing and thawing) Shipping: Blue Ice

IMAGES




FCM/FACS (Flow Cytometry)

Figure 3. Affinity ranking of different Rabbit anti-CD38 mAb clones by titration of different concentration onto Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.



ELISA

Figure 3. Affinity ranking of different Rabbit anti-CD38 mAb clones by titration of different concentration onto Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.



ELISA

Figure 3. Affinity ranking of different Rabbit anti-CD38 mAb clones by titration of different concentration onto Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.

ADDITIONAL INFORMATION

Related Product Information for anti-CD38 antibody	CD antigen CD38 is also known as ADP-ribosyl cyclase 1, which belongs to the ADP-ribosyl cyclase family. CD38 is expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma. CD38 is a multifunctional ectoenzyme that catalyzes the synthesis and hydrolysis of cyclic ADP-ribose (cADPR) from NAD ⁺ to ADP-ribose. These reaction products are essential for the regulation of intracellular Ca ²⁺ . The loss of CD38 function is associated with impaired immune responses, metabolic disturbances, and behavioral modifications. The CD38 protein is a marker of cell activation. It has been connected to HIV infection, leukemias, myelomas, solid tumors, type II diabetes mellitus and bone metabolism. CD38 has been used as a prognostic marker in leukemia.
Product Categories/Family for anti-CD38 antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	55977782
NCBI GeneID	952
UniProt Accession #	P28907; O00121; O00122; Q96HY4
Molecular Weight	300
NCBI Official Full Name	ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1
NCBI Official Synonym Full Names	CD38 molecule
NCBI Official Symbol	CD38
NCBI Official Synonym Symbols	T10; ADPRC 1
NCBI Protein Information	ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1; cADPr hydrolase 1; CD38 antigen (p45); NAD(+) nucleosidase; ADP-ribosyl cyclase 1; 2'-phospho-ADP-ribosyl cyclase; 2'-phospho-cyclic-ADP-ribose transferase; 2'-phospho-ADP-ribosyl cyclase/2'-phospho-cycli
UniProt Protein Name	ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1
UniProt Gene Name	CD38
UniProt Synonym Gene Names	ADPRC 1; cADPr hydrolase 1
UniProt Entry Name	CD38_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

