

Rabbit anti-Human CTLA4 Monoclonal Antibody | anti-CTLA4 antibody

Anti-CTLA4 antibody (DM51), Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA47290	Rabbit	Human	Flow Cytometry, Functional Assay, ELISA


PRODUCT SYNONYMS

CTLA4; Monoclonal Antibody; Anti-CTLA4 antibody (DM51); Rabbit mAb; CD152; anti-CTLA4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	DM51
Purity/Purification	Purified from cell culture supernatant by affinity chromatography
Form/Format	Liquid. 50% Glycerol; PBS, pH7.4; 0.1% BSA; 0.1% Procline 300
Applicable Applications for anti-CTLA4 antibody	FCM/FACS (Flow Cytometry), ELISA
Immunogen	Recombinant human CTLA4 (Gln36-Ser161) produced by using human HEK293 cells
Reconstitution	Reconstitute with deionized water
Preparation and Storage	Store at -20 degree C for 12 months (Avoid repeated freezing and thawing)

IMAGES




FCM/FACS (Flow Cytometry)

Figure 1. ELISA plate pre-coated by 2 ug/ml (100 ul/well) Human CTLA-4 protein, mFc-His tagged protein (PME100017) can bind Rabbit anti-CTLA-4 monoclonal antibody(clone: DM50) in a linear range of 1-100 ng/ml.



FCM/FACS (Flow Cytometry)

Figure 1. ELISA plate pre-coated by 2 ug/ml (100 ul/well) Human CTLA-4 protein, mFc-His tagged protein (PME100017) can bind Rabbit anti-CTLA-4 monoclonal antibody(clone: DM50) in a linear range of 1-100 ng/ml.



ELISA

Figure 1. ELISA plate pre-coated by 2 ug/ml (100 ul/well) Human CTLA-4 protein, mFc-His tagged protein (PME100017) can bind Rabbit anti-CTLA-4 monoclonal antibody(clone: DM50) in a linear range of 1-100 ng/ml.

ADDITIONAL INFORMATION

Related Product Information for anti-CTLA4 antibody

Leukocyte surface antigen CD47 is also known as Antigenic surface determinant protein OA3, Integrin-associated protein (IAP) and Protein MER6. CD47 contains 1 Ig-like V-type (immunoglobulin-like) domain. CD47 is very broadly distributed on normal adult tissues. CD47 has a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins and plays an important role in memory formation and synaptic plasticity in the hippocampus by similarity. CD47 is the receptor for SIRPA, binding to which prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. CD47 Interaction with SIRPG mediates cell-cell adhesion, enhances superantigen-dependent T-cell-mediated proliferation and costimulates T-cell activation.

NCBI AND UNIPROT INFORMATION

NCBI GI #	27735177
NCBI GeneID	1493
UniProt Accession #	P16410; Q95653; Q0PP65; Q52MC1; Q53TD5; Q5S005; Q8WXJ1; Q96P43; Q9UKN9; A0N1S0; E9PDH0
Molecular Weight	24,656 Da
NCBI Official Full Name	Cytotoxic T-lymphocyte protein 4
NCBI Official Synonym Full Names	cytotoxic T-lymphocyte-associated protein 4
NCBI Official Symbol	CTLA4
NCBI Official Synonym Symbols	CD; GSE; GRD4; CD152; CTLA-4; IDDM12; CELIAC3
NCBI Protein Information	cytotoxic T-lymphocyte protein 4; CD152 isoform; celiac disease 3; cytotoxic T-lymphocyte antigen 4; insulin-dependent diabetes mellitus 12; cytotoxic T-lymphocyte-associated antigen 4; cytotoxic T-lymphocyte-associated serine esterase-4; cytotoxic T lymph
UniProt Protein Name	Cytotoxic T-lymphocyte protein 4
UniProt Gene Name	CTLA4
UniProt Synonym Gene Names	CD152; CTLA-4
UniProt Entry Name	CTLA4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/ctla4-antibody-clone-dm51-flow-cytometry-elisa/47290>