

Rabbit anti-Human 41BB Ligand Monoclonal Antibody | anti-41BB antibody

Anti-41BB Ligand antibody (DM68), Rabbit mAb

CATALOG # AAA47300	HOST Rabbit	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay, ELISA
------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

41BB Ligand; Monoclonal Antibody; Anti-41BB Ligand antibody (DM68); Rabbit mAb; 4-1BB Ligand; TNFSF9; CD137L; anti-41BB antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	DM68
Purity/Purification	Purified from cell culture supernatant by affinity chromatography
Form/Format	Liquid. 50% Glycerol; PBS, pH7.4; 0.1% BSA; 0.1% Procline 300
Applicable Applications for anti-41BB antibody	FCM/FACS (Flow Cytometry), ELISA
Immunogen	Recombinant human 4-1BB Ligand (Pro52-Glu254) produced by using human HEK293 cells
Reconstitution	Reconstitute with deionized water
Preparation and Storage	Store at -20 degree C for 12 months (Avoid repeated freezing and thawing)

IMAGES




FCM/FACS (Flow Cytometry)

Figure 3. Flow cytometry data of serially titrated Rabbit anti-4-1BB Ligand monoclonal antibody (clone: DM68) on Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.



FCM/FACS (Flow Cytometry)

Figure 3. Flow cytometry data of serially titrated Rabbit anti-4-1BB Ligand monoclonal antibody (clone: DM68) on Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.



ELISA

Figure 3. Flow cytometry data of serially titrated Rabbit anti-4-1BB Ligand monoclonal antibody (clone: DM68) on Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.

ADDITIONAL INFORMATION

Related Product Information for anti-41BB antibody	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This transmembrane cytokine is a bidirectional signal transducer that acts as a ligand for TNFRSF9/4-1BB, which is a costimulatory receptor molecule in T lymphocytes. This cytokine and its receptor are involved in the antigen presentation process and in the generation of cytotoxic T cells. The receptor TNFRSF9/4-1BB is absent from resting T lymphocytes but rapidly expressed upon antigenic stimulation. The ligand encoded by this gene, TNFSF9/4-1BBL, has been shown to reactivate anergic T lymphocytes in addition to promoting T lymphocyte proliferation. This cytokine has also been shown to be required for the optimal CD8 responses in CD8 T cells. This cytokine is expressed in carcinoma cell lines, and is thought to be involved in T cell-tumor cell interaction.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507609
NCBI GeneID	8744
NCBI Accession #	NP_003802.1
NCBI GenBank Nucleotide #	NM_003811.3
UniProt Accession #	P41273; Q2M3S2
Molecular Weight	26,625 Da
NCBI Official Full Name	tumor necrosis factor ligand superfamily member 9
NCBI Official Synonym Full Names	tumor necrosis factor (ligand) superfamily, member 9
NCBI Official Symbol	TNFSF9
NCBI Official Synonym Symbols	CD137L; 4-1BB-L
NCBI Protein Information	tumor necrosis factor ligand superfamily member 9; 4-1BBL; 4-1BB ligand; receptor 4-1BB ligand; homolog of mouse 4-1BB-L
UniProt Protein Name	Tumor necrosis factor ligand superfamily member 9
UniProt Gene Name	TNFSF9
UniProt Synonym Gene Names	4-1BBL
UniProt Entry Name	TNFL9_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/41bb-ligand-antibody-clone-dm68-flow-cytometry-elisa/47300>