

Rabbit anti-Human B7-2 Monoclonal Antibody | anti-B7-2 antibody

Anti-B7-2 antibody (DM85), Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA47317	Rabbit	Human	Flow Cytometry, Functional Assay, ELISA


PRODUCT SYNONYMS

B7-2; Monoclonal Antibody; Anti-B7-2 antibody (DM85); Rabbit mAb; CD86; B70; CD28LG2; LAB72; MGC34413; anti-B7-2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	DM85
Purity/Purification	Purified from cell culture supernatant by affinity chromatography
Form/Format	Liquid. 50% Glycerol; PBS, pH7.4; 0.1% BSA; 0.1% Procline 300
Applicable Applications for anti-B7-2 antibody	FCM/FACS (Flow Cytometry), ELISA
Immunogen	Recombinant human B7-2 (Leu26-Pro247) produced by using human HEK293 cells
Reconstitution	Reconstitute with deionized water
Preparation and Storage	Store at -20 degree C for 12 months (Avoid repeated freezing and thawing)

IMAGES



FCM/FACS (Flow Cytometry)

Figure 1. ELISA plate pre-coated by 2 ug/ml (100 ul/well) Human B7-2 protein, mFc-His tagged protein PME100034 can bind Rabbit anti-B7-2 monoclonal antibody (clone: DM85) in a linear range of 1-25 ng/ml.



ELISA

Figure 1. ELISA plate pre-coated by 2 ug/ml (100 ul/well) Human B7-2 protein, mFc-His tagged protein PME100034 can bind Rabbit anti-B7-2 monoclonal antibody (clone: DM85) in a linear range of 1-25 ng/ml.

ADDITIONAL INFORMATION

Related Product Information for anti-B7-2 antibody	This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript variants encoding different isoforms.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	317373339
NCBI GeneID	942
UniProt Accession #	P42081; Q13655; Q6FHB1; Q6GTS4; Q7M4L5; A0N0P0; B7Z2F3; B7Z702; E7ETN5; E9PC27
Molecular Weight	329
NCBI Official Full Name	T-lymphocyte activation antigen CD86
NCBI Official Synonym Full Names	CD86 molecule
NCBI Official Symbol	CD86
NCBI Official Synonym Symbols	B70; B7-2; B7.2; LAB72; CD28LG2
NCBI Protein Information	T-lymphocyte activation antigen CD86; BU63; FUN-1; CTLA-4 counter-receptor B7.2; B-lymphocyte activation antigen B7-2; CD86 antigen (CD28 antigen ligand 2, B7-2 antigen)
UniProt Protein Name	T-lymphocyte activation antigen CD86
UniProt Gene Name	CD86
UniProt Synonym Gene Names	CD28LG2
UniProt Entry Name	CD86_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/b7-2-antibody-clone-dm85-flow-cytometry-elisa/47317>