

CD56 recombinant protein

Human CD56 Protein, His Tag

CATALOG #
AAA47347

HOST
HEK293 cells


PRODUCT SYNONYMS

CD56; Human CD56 Protein; His Tag; NCAM1; MSK39; NCAM; CD56 recombinant protein

PRODUCT SPECIFICATIONS


Host	HEK293 cells
Purity/Purification	>90%. Affinity purification
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose is added as protectants before lyophilization.
Sequence	CD56(Leu22 0-Gly718)+ 6×Histag
Origin	Human
Reconstitution	Reconstitute with deionized water
Tag	C-6×His tag
Preparation and Storage	Store at -80 degree C for 12 months (Avoid repeated freezing and thawing)

IMAGES



SDS-PAGE

Figure 1. Human CD56 Protein, His Tag on SDS-PAGE under reducing condition.



ELISA

Figure 1. Human CD56 Protein, His Tag on SDS-PAGE under reducing condition.

ADDITIONAL INFORMATION

Related Product Information for CD56 recombinant protein	This gene encodes a cell adhesion protein which is a member of the immunoglobulin superfamily. The encoded protein is involved in cell-to-cell interactions as well as cell-matrix interactions during development and differentiation. The encoded protein plays a role in the development of the nervous system by regulating neurogenesis, neurite outgrowth, and cell migration. This protein is also involved in the expansion of T lymphocytes, B lymphocytes and natural killer (NK) cells which play an important role in immune surveillance. This protein plays a role in signal transduction by interacting with fibroblast growth factor receptors, N-cadherin and other components of the extracellular matrix and by triggering signalling cascades involving FYN-focal adhesion kinase (FAK), mitogen-activated protein kinase (MAPK), and phosphatidylinositol 3-kinase (PI3K). One prominent isoform of this gene, cell surface molecule CD56, plays a role in several myeloproliferative disorders such as acute myeloid leukemia and differential expression of this gene is associated with differential disease progression. For example, increased expression of CD56 is correlated with lower survival in acute myeloid leukemia patients whereas increased severity of COVID-19 is correlated with decreased abundance of CD56-expressing NK cells in peripheral blood. Alternative splicing results in multiple transcript variants encoding distinct protein isoforms.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	205830665
NCBI GeneID	4684
UniProt Accession #	P13591; P13592; P13593; Q05C58; Q15829; Q16180; Q16209; Q59FL7; Q86X47; Q96CJ3; A8K8T8
Molecular Weight	858
NCBI Official Full Name	Neural cell adhesion molecule 1
NCBI Official Synonym Full Names	neural cell adhesion molecule 1
NCBI Official Symbol	NCAM1
NCBI Official Synonym Symbols	CD56; NCAM; MSK39
NCBI Protein Information	neural cell adhesion molecule 1; neural cell adhesion molecule, NCAM; antigen recognized by monoclonal antibody 5.1H11
UniProt Protein Name	Neural cell adhesion molecule 1
UniProt Gene Name	NCAM1
UniProt Synonym Gene Names	NCAM; N-CAM-1; NCAM-1
UniProt Entry Name	NCAM1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/ncam1-recombinant-protein-hek293-cells/47347>