

VEGF recombinant protein

Human VEGF Protein, hFc Tag

CATALOG #
AAA47385

HOST
HEK293 cells


PRODUCT SYNONYMS

VEGF; Human VEGF Protein; hFc Tag; MVCD1; VPF; VEGF recombinant protein

PRODUCT SPECIFICATIONS


Host	HEK293 cells
Purity/Purification	Affinity purification
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose is added as protectants before lyophilization.
Sequence	VEGF(Ala27 -Arg232)+h Fc(Glu99-A la330)
Origin	Human
Reconstitution	Reconstitute with deionized water
Tag	C-Human Fc
Preparation and Storage	Store at -80 degree C for 12 months (Avoid repeated freezing and thawing)

IMAGES



SDS-PAGE

Figure 1. Human VEGFA Protein, hFc Tag on SDS-PAGE under reducing conditions.



ELISA

Figure 1. Human VEGFA Protein, hFc Tag on SDS-PAGE under reducing conditions.

ADDITIONAL INFORMATION

Related Product Information for VEGF recombinant protein	This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site. [provided by RefSeq, Nov 2015]
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	324120930
NCBI GeneID	7422
NCBI Accession #	NP_001191313.1
NCBI GenBank Nucleotide #	NM_001204384.1
UniProt Accession #	P15692; O60720; O75875; Q074Z4; Q16889; B5BU86; H0Y2S8; H0Y407; H0Y414; H0Y462; H0Y8N2; H3BLW7
Molecular Weight	27,042 Da
NCBI Official Full Name	vascular endothelial growth factor A isoform q
NCBI Official Synonym Full Names	vascular endothelial growth factor A
NCBI Official Symbol	VEGFA
NCBI Official Synonym Symbols	VPF; VEGF; MVCD1
NCBI Protein Information	vascular endothelial growth factor A; vascular permeability factor
UniProt Protein Name	Vascular endothelial growth factor A
UniProt Gene Name	VEGFA
UniProt Synonym Gene Names	VEGF; VEGF-A; VPF
UniProt Entry Name	VEGFA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.