

Rabbit anti-Human 4-1BB Monoclonal Antibody | anti-4-1BB antibody

Anti-4-1BB antibody (DM67), Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA47415	Rabbit	Human	Flow Cytometry, Functional Assay, ELISA


PRODUCT SYNONYMS

4-1BB; Monoclonal Antibody; Anti-4-1BB antibody (DM67); Rabbit mAb; TNFRSF9; CD137; CDw137; ILA; anti-4-1BB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	DM67
Purity/Purification	Purified from cell culture supernatant by affinity chromatography
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose is added as protectants before lyophilization. 0.1% Procline 300
Applicable Applications for anti-4-1BB antibody	FCM/FACS (Flow Cytometry), ELISA
Immunogen	Recombinant human 4-1BB (Leu24-Gln186) produced by using human HEK293 cells
Reconstitution	Reconstitute with deionized water
Preparation and Storage	Store at -20 degree C for 12 months (Avoid repeated freezing and thawing)

IMAGES




FCM/FACS (Flow Cytometry)

Figure 3. Flow cytometry data of serially titrated Rabbit anti-4-1BB monoclonal antibody (clone: DM67) on Expi 293 cell line transfected with human 4-1BB. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.



FCM/FACS (Flow Cytometry)

Figure 3. Flow cytometry data of serially titrated Rabbit anti-4-1BB monoclonal antibody (clone: DM67) on Expi 293 cell line transfected with human 4-1BB. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.



ELISA

Figure 3. Flow cytometry data of serially titrated Rabbit anti-4-1BB monoclonal antibody (clone: DM67) on Expi 293 cell line transfected with human 4-1BB. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.

ADDITIONAL INFORMATION

Related Product Information for anti-4-1BB antibody

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contributes to the clonal expansion, survival, and development of T cells. It can also induce proliferation in peripheral monocytes, enhance T cell apoptosis induced by TCR/CD3 triggered activation, and regulate CD28 co-stimulation to promote Th1 cell responses. The expression of this receptor is induced by lymphocyte activation. TRAF adaptor proteins have been shown to bind to this receptor and transduce the signals leading to activation of NF-kappaB.

NCBI AND UNIPROT INFORMATION

NCBI GI #	5730095
NCBI GeneID	3604
NCBI Accession #	NP_001552.2
NCBI GenBank Nucleotide #	NM_001561.5
UniProt Accession #	Q07011; AAB34924
Molecular Weight	27,899 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 9
NCBI Official Synonym Full Names	tumor necrosis factor receptor superfamily, member 9
NCBI Official Symbol	TNFRSF9
NCBI Official Synonym Symbols	ILA; 4-1BB; CD137; CDw137
NCBI Protein Information	tumor necrosis factor receptor superfamily member 9; CD137 antigen; T cell antigen ILA; T-cell antigen ILA; 4-1BB ligand receptor; homolog of mouse 4-1BB; receptor protein 4-1BB; T-cell antigen 4-1BB homolog; induced by lymphocyte activation (ILA); interl
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 9
UniProt Gene Name	TNFRSF9
UniProt Synonym Gene Names	CD137; ILA
UniProt Entry Name	TNR9_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/4-1bb-antibody-clone-dm67-flow-cytometry-elisa/47415>