

Rabbit anti-Human B7-H2 Monoclonal Antibody | anti-B7-H2 antibody

Anti-B7-H2 antibody (DM98), Rabbit mAb

CATALOG # AAA47423	HOST Rabbit	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay, ELISA
------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

B7-H2; Monoclonal Antibody; Anti-B7-H2 antibody (DM98); Rabbit mAb; ICOSLG; B7H2; B7RP-1; B7RP1; CD275; GL50; ICOS-L; ICOSL; LICOS; ICOS ligand; anti-B7-H2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	DM98
Purity/Purification	Purified from cell culture supernatant by affinity chromatography
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose is added as protectants before lyophilization. 0.1% Procline 300
Applicable Applications for anti-B7-H2 antibody	FCM/FACS (Flow Cytometry), ELISA
Immunogen	Recombinant human B7-H2 (Asp19-Thr256) produced by using human HEK293 cells
Reconstitution	Reconstitute with deionized water
Preparation and Storage	Store at -20 degree C for 12 months (Avoid repeated freezing and thawing)

IMAGES



FCM/FACS (Flow Cytometry)

Figure 1. ELISA plate pre-coated by 2 ug/ml (100 ul/well) Human B7-H2 protein, mFc-His tagged protein PME100029 can bind Rabbit anti-B7-H2 monoclonal antibody (clone: DM98) in a linear range of 3.2-80 ng/ml.



ELISA

Figure 1. ELISA plate pre-coated by 2 ug/ml (100 ul/well) Human B7-H2 protein, mFc-His tagged protein PME100029 can bind Rabbit anti-B7-H2 monoclonal antibody (clone: DM98) in a linear range of 3.2-80 ng/ml.

ADDITIONAL INFORMATION

Related Product Information for anti-B7-H2 antibody	Inducible co-stimulator ligand (ICOSL), also known as B7-H2, is a member of the B7 family of co-stimulatory molecules related to B7-1 and B7-2. The protein is the ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion; induces also B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-stimulating memory T-cell function.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	19855066
NCBI GeneID	102723996
UniProt Accession #	O75144; Q9HD18; Q9NRQ1; A8MUZ1
Molecular Weight	20,120 Da
NCBI Official Full Name	ICOS ligand
NCBI Official Symbol	LOC102723996
NCBI Protein Information	ICOS ligand
UniProt Protein Name	ICOS ligand
UniProt Gene Name	ICOSLG
UniProt Synonym Gene Names	B7H2; B7RP1; ICOSL; KIAA0653; B7-H2; B7RP-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/b7-h2-antibody-clone-dm98-flow-cytometry-elisa/47423>