

Rabbit anti-Human PD-L1 Monoclonal Antibody | anti-PD-L1 antibody

Anti-PD-L1 antibody (DM124), Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA47448	Rabbit	Human	Flow Cytometry, Functional Assay, ELISA


PRODUCT SYNONYMS

PD-L1; Monoclonal Antibody; Anti-PD-L1 antibody (DM124); Rabbit mAb; CD274; B7-H1; PDCD1L1; PDCD1LG1; anti-PD-L1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	DM124
Purity/Purification	Purified from cell culture supernatant by affinity chromatography
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose is added as protectants before lyophilization. 0.1% Procline 300
Applicable Applications for anti-PD-L1 antibody	FCM/FACS (Flow Cytometry), ELISA
Immunogen	Recombinant human PD-L1 (Phe19-Arg328) produced by using human HEK293 cells
Reconstitution	Reconstitute with deionized water
Preparation and Storage	Store at -20 degree C for 12 months (Avoid repeated freezing and thawing)

IMAGES



FCM/FACS (Flow Cytometry)

Figure 1. ELISA plate pre-coated by 1 ug/ml (100 ul/well) Human PD-L1 protein, mFc His tagged protein PME100023 can bind Rabbit anti-PD-L1 monoclonal antibody (clone: DM124) in a linear range of 0.1-15 ng/ml.



ELISA

Figure 1. ELISA plate pre-coated by 1 ug/ml (100 ul/well) Human PD-L1 protein, mFc His tagged protein PME100023 can bind Rabbit anti-PD-L1 monoclonal antibody (clone: DM124) in a linear range of 0.1-15 ng/ml.

ADDITIONAL INFORMATION

Related Product Information for anti-PD-L1 antibody	This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	390979639
NCBI GeneID	29126
NCBI Accession #	NP_001254635.1
NCBI GenBank Nucleotide #	NM_001267706.1
UniProt Accession #	Q9NZQ7; Q14CJ2; Q2V8D5; Q66RK1; Q6WEX4; Q9NUZ5; B2RBA2; B4DU27
Molecular Weight	33,275 Da
NCBI Official Full Name	programmed cell death 1 ligand 1 isoform b
NCBI Official Synonym Full Names	CD274 molecule
NCBI Official Symbol	CD274
NCBI Official Synonym Symbols	B7-H; B7H1; PDL1; PD-L1; PDCD1L1; PDCD1LG1
NCBI Protein Information	programmed cell death 1 ligand 1; B7 homolog 1; CD274 antigen; PDCD1 ligand 1; programmed death ligand 1
UniProt Protein Name	Programmed cell death 1 ligand 1
UniProt Gene Name	CD274
UniProt Synonym Gene Names	B7H1; PDCD1L1; PDCD1LG1; PDL1; PD-L1; PDCD1 ligand 1; Programmed death ligand 1; B7-H1
UniProt Entry Name	PD1L1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/pd-l1-antibody-clone-dm124-flow-cytometry-elisa/47448>