

CD142 recombinant protein

Human CD142 Protein, hFc Tag

CATALOG #
AAA47552

HOST
HEK293

PRODUCT SYNONYMS

CD142; Human CD142 Protein; hFc Tag; TF; Coagulation factor III; F3; CD142 recombinant protein

PRODUCT SPECIFICATIONS

Host	HEK293
Purity/Purification	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
Tag	C-Human Fc tag
Preparation and Storage	Store at -20 degree C to -80 degree C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80 degree C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

IMAGES



SDS-PAGE

Figure 1. Human CD142 Protein, hFc Tag on SDS-PAGE under reducing condition.



ELISA

Figure 1. Human CD142 Protein, hFc Tag on SDS-PAGE under reducing condition.

ADDITIONAL INFORMATION

Related Product Information for CD142 recombinant protein	Recombinant human CD142 protein with C-terminal human Fc tag This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces, for example, on monocytes. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. Platelets and monocytes have been shown to express this coagulation factor under procoagulatory and proinflammatory stimuli, and a major role in HIV-associated coagulopathy has been described. Platelet-dependent monocyte expression of coagulation factor III has been described to be associated with Coronavirus Disease 2019 (COVID-19) severity and mortality. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Aug 2020]
Product Categories/Family for CD142 recombinant protein	Bioactive Protein

NCBI AND UNIPROT INFORMATION

NCBI GI #	296010912
NCBI GeneID	2152
NCBI Accession #	NP_001171567.1
NCBI GenBank Nucleotide #	NM_001178096.1
UniProt Accession #	P13726; Q6FHG2; Q86WH4; D3DT47
Molecular Weight	33,068 Da
NCBI Official Full Name	tissue factor isoform 2
NCBI Official Synonym Full Names	coagulation factor III (thromboplastin, tissue factor)
NCBI Official Symbol	F3
NCBI Official Synonym Symbols	TF; TFA; CD142
NCBI Protein Information	tissue factor
UniProt Protein Name	Tissue factor
UniProt Gene Name	F3
UniProt Synonym Gene Names	TF
UniProt Entry Name	TF_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/f3-recombinant-protein-hek293/47552>