

CCR8 protein

Human CCR8 full length protein-synthetic nanodisc

CATALOG #
AAA47619

HOST
HEK293



PRODUCT SYNONYMS

CCR8; Human CCR8 full length protein-synthetic nanodisc; CC-CKR-8; CCR-8; CDw198; CKRL1; CMKBR8; CMKBRL2; CY6; GPRCY6; TER1; CCR8 protein

PRODUCT SPECIFICATIONS

Host	HEK293
Form/Format	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
Preparation and Storage	Store at -20 degree C to -80 degree C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80 degree C (Avoid repeated freezing and thawing).

IMAGES



Application Data

Figure 3. Human CCR8-Nanodisc can bind Anti-CCR8 antibody (BME100115) with an affinity constant of 1.408 nM as determined in a SPR assay.

WB (Western Blot)

Figure 3. Human CCR8-Nanodisc can bind Anti-CCR8 antibody (BME100115) with an affinity constant of 1.408 nM as determined in a SPR assay.

ADDITIONAL INFORMATION

Related Product Information for CCR8 protein

A member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptors are important for the migration of various cell types into the inflammatory sites. This receptor protein preferentially expresses in the thymus. I-309, thymus activation-regulated cytokine (TARC) and macrophage inflammatory protein-1 beta (MIP-1 beta) have been identified as ligands of this receptor. Studies of this receptor and its ligands suggested its role in regulation of monocyte chemotaxis and thymic cell apoptosis. More specifically, this receptor may contribute to the proper positioning of activated T cells within the antigenic challenge sites and specialized areas of lymphoid tissues. This gene is located at the chemokine receptor gene cluster region.

Product Categories/Family for CCR8 protein

FLP 489; Full Length Transmembrane Proteins

NCBI AND UNIPROT INFORMATION

NCBI GI #	1707884
NCBI GeneID	1237
UniProt Accession #	P51685; Q3KNQ8; Q3KNR3; Q9BYX5; B2RC64
NCBI Official Full Name	C-C chemokine receptor type 8
NCBI Official Synonym Full Names	chemokine (C-C motif) receptor 8
NCBI Official Symbol	CCR8
NCBI Official Synonym Symbols	CY6; TER1; CCR-8; CKRL1; CDw198; CMKBR8; GPRCY6; CMKBRL2; CC-CKR-8
NCBI Protein Information	C-C chemokine receptor type 8; CC chemokine receptor 8; chemokine receptor-like 1; chemokine (C-C) receptor 8; CC chemokine receptor CHEMR1; CC-chemokine receptor chemr1; chemokine (C-C) receptor-like 2
UniProt Protein Name	C-C chemokine receptor type 8
UniProt Gene Name	CCR8
UniProt Synonym Gene Names	CKRL1; CMKBR8; CMKBRL2; C-C CKR-8; CC-CKR-8; CCR-8; CKR-L1; GPRCY6
UniProt Entry Name	CCR8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/hek293-protein/ccr8/47619>