

Goat anti Rabbit IgG, Fc Fragment Specific Antibody, Peroxidase Conjugated Secondary Antibody

Goat anti Rabbit IgG, Fc Fragment Specific Antibody, Peroxidase Conjugated

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA47968	Goat	Rabbit	Immunocytochemistry, Immunohistochemistry, Western Blot, ELISA

PRODUCT SYNONYMS

Goat anti Rabbit IgG; Fc Fragment Specific Antibody; Peroxidase Conjugated; Peroxidase Conjugated secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Rabbit
Isotype	IgG
Specificity	By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG Fc fragment. This antibody may cross-react with IgG of other species.
Purity/Purification	Purified by peptide immuno-affinity chromatography
Form/Format	Liquid; Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.05% Pro-Clean 400
Concentration	1.0 mg/ml (varies by lot)
Applicable Applications for Goat anti Rabbit IgG, Fc Fragment Specific Antibody, Peroxidase Conjugated secondary antibody	ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Rabbit IgG Fc fragment
Preparation and Storage	Store at 2 to 8°C.

ADDITIONAL INFORMATION

Related Product Information for Goat anti Rabbit IgG, Fc Fragment Specific Antibody, Peroxidase Conjugated secondary antibody	IgG is the most abundant immunoglobulin in blood and plasma, playing a crucial role in immunity. It functions by neutralizing toxins, activating the complement system, immobilizing pathogens, and initiating opsonization and cell-mediated cytotoxicity. This goat anti-IgG Fc specific polyclonal antibody is specifically designed to bind only to the Fc portion of the IgG molecule, ensuring no cross-reactivity with the Fab fragments or light chains
Product Categories/Family for Goat anti Rabbit IgG, Fc Fragment Specific Antibody, Peroxidase Conjugated secondary antibody	Secondary Antibodies

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/goat-rabbit-secondary-antibody/goat-anti-rabbit-igg-fc-fragment-specific-antibody-peroxidase-conjugated/47968>