

Mouse CD55 Monoclonal Antibody | anti-CD55 antibody

MOUSE ANTI HUMAN CD55


CATALOG # AAA49279	HOST Mouse	APPLICATIONS Western Blot, Flow Cytometry, Functional Assay, Immunohistochemistry
------------------------------	----------------------	---

PRODUCT SYNONYMS


CD55; Monoclonal Antibody; MOUSE ANTI HUMAN CD55; anti-CD55 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Number	67
Form/Format	Purified Purified IgG - liquid
Concentration	IgG concentration 1.0 mg/ml (varies by lot)
Sequence Length	551
Applicable Applications for anti-CD55 antibody	WB (Western Blot), FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry)
Perservative Stabilisers	0.09% Sodium Azide Preparation
Immunogen	K562 cells Histology Positive Control Tissue
Target Species	Human
Preparation and Storage	Store at 4 degree C or at -20 degree C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life: 18 months from date of despatch.



Application Data
Staining of human peripheral blood lymphocytes with Mouse anti Human CD55



Application Data
Staining of human peripheral blood monocytes with Mouse anti Human CD55:RPE



Application Data
Staining of human peripheral blood monocytes with Mouse anti Human CD55

ADDITIONAL INFORMATION

Related Product Information for anti-CD55 antibody

Mouse anti Human CD55 antibody, clone 67 recognizes the human CD55 cell surface antigen, a GPI linked molecule also known as decay accelerating factor (DAF). CD55 is expressed by a wide range of cell types. Erythrocyte expression is very weak. CD55 is the complement regulatory protein, decay accelerating factor (DAF) (Lublin and Atkinson 1989). Human CD55 is a ~70kDa glycoprotein (in erythrocytes) anchored in the membrane by glycosylphosphatidylinositol tail. In other cells the apparent molecular weight is somewhat larger. It has a substantial content of O-glycans, and also on N-glycan. DAF binds to activated C4b or C3b complement fragments on the cell surface, preventing the assembly and accelerating the decay of both classical and alternative pathways. DAF carries the Cromer related blood group antigens. DAF has a wide distribution on cells in non-haemopoietic tissues, particularly epithelium and is specifically found at the foetal-maternal interface in placenta (Holmes et al. 1990 and Yang et al. 2009). Soluble forms of DAF are found, for example, in plasma, saliva and urine. The antigen on erythrocytes is pronase and chymotrypsin sensitive, but resistant to trypsin.

NCBI AND UNIPROT INFORMATION

NCBI GI #	10835143
NCBI GeneID	1604
NCBI Accession #	NP_000565.1
NCBI GenBank Nucleotide #	NM_000574.4
UniProt Accession #	P08174; P09679; P78361; Q14UF2; Q14UF3; Q14UF4; Q14UF5; Q14UF6; B1AP14; D3DT83; D3DT84; E7ER69
Molecular Weight	59,038 Da

NCBI Official Full Name	complement decay-accelerating factor isoform 1 preproprotein
NCBI Official Synonym Full Names	CD55 molecule, decay accelerating factor for complement (Cromer blood group)
NCBI Official Symbol	CD55
NCBI Official Synonym Symbols	CR; TC; DAF; CROM
NCBI Protein Information	complement decay-accelerating factor; CD55 antigen
UniProt Protein Name	Complement decay-accelerating factor
UniProt Gene Name	CD55
UniProt Synonym Gene Names	CR; DAF
UniProt Entry Name	DAF_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd55-antibody-clone-67-western-blot-flow-cytometry-immunohistochemistry-elisa/49279>