

Mouse CD25 Monoclonal Antibody | anti-CD25 antibody

MOUSE ANTI HUMAN CD25


CATALOG #	HOST	APPLICATIONS
AAA49334	Mouse	Flow Cytometry, Functional Assay

PRODUCT SYNONYMS


CD25; Monoclonal Antibody; MOUSE ANTI HUMAN CD25; anti-CD25 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Number	MEM-181
Form/Format	Purified Purified IgG - liquid
Concentration	IgG concentration 1 mg/ml (varies by lot)
Sequence Length	272
Applicable Applications for anti-CD25 antibody	FCM/FACS (Flow Cytometry)
Perservative Stabilisers	0.09% Sodium Azide (NaN_3)
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Target Species	Human
Immunogen	Human PHA blasts; day 3 of culture
Preparation and Storage	It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquot at $2-8^{\circ}$ for short term use (up to 4 weeks) and store the remaining aliquots at -20°C Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended




Application Data
Staining of PHA activated human peripheral blood lymphocytes with Mouse anti Human CD25



Application Data
Staining of PHA stimulated human peripheral blood lymphocytes with Mouse anti Human CD25: RPE- Alexa Fluor 488 (AAA49334A488)



Application Data
Staining of PHA stimulated human peripheral blood lymphocytes with Mouse anti Human CD25:FITC



Application Data
Staining of PHA stimulated human lymphocyte cells with Mouse anti Human CD25

ADDITIONAL INFORMATION

Related Product Information for anti-CD25 antibody	Mouse anti Human CD25, clone MEM-181 recognizes the 55kD alpha subunit of the human IL-2 receptor, also known as p55 or TAC antigen, CD25 is a type 1 transmembrane protein with two Sushi domains, also known as short consensus repeats (SCRs) or complement control protein (CCP) modules (Norman et al. 1991) located within its extracellular domain. The IL-2 receptor exists in three forms. A high affinity form consisting of a non-covalently linked heterodimer composed of the alpha subunit (CD25) and the IL-2 receptor beta subunit also known as CD122 or p75, a medium affinity beta subunit (CD122) monomer or a low affinity alpha (CD25) subunit monomer. CD25 is expressed by activated T lymphocytes and activated B lymphocytes responding to antigen or mitogen stimulation. CD25 is also expressed in some thymocytes and oligodendrocytes. In disease, elevated expression of CD25 is noted in a number of chronic inflammatory conditions, tuberculoid leprosy patients demonstrate markedly elevated levels of circulating CD25high FoxP3+ regulatory T cells (T-regs) (Attia et al. 2010). Elevated levels of CD25 antigen expression are often seen in cases of non-Hodgkin 's lymphoma and diffuse large B cell lymphoma (Fujiwara et al.2013).
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557667
NCBI GeneID	3559
NCBI Accession #	NP_000408.1
NCBI GenBank Nucleotide #	NM_000417.2
UniProt Accession #	P01589; Q5W007

NCBI Official Full Name	interleukin-2 receptor subunit alpha
NCBI Official Synonym Full Names	interleukin 2 receptor, alpha
NCBI Official Symbol	IL2RA
NCBI Official Synonym Symbols	p55; CD25; IL2R; TCGFR; IDDM10
NCBI Protein Information	interleukin-2 receptor subunit alpha; IL-2 receptor subunit alpha; IL-2R subunit alpha; TAC antigen
UniProt Protein Name	Interleukin-2 receptor subunit alpha
UniProt Gene Name	IL2RA
UniProt Synonym Gene Names	IL-2 receptor subunit alpha; IL-2-RA; IL-2R subunit alpha; IL2-RA
UniProt Entry Name	IL2RA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd25-antibody-clone-mem-181-flow-cytometry-elisa/49334>