

Mouse CD63 Monoclonal Antibody | anti-CD63 antibody

MOUSE ANTI HUMAN CD63

CATALOG # AAA49336	HOST Mouse	APPLICATIONS Immunoprecipitation, Immunofluorescence, Flow Cytometry, Functional Assay
------------------------------	----------------------	--

PRODUCT SYNONYMS

CD63; Monoclonal Antibody; MOUSE ANTI HUMAN CD63; anti-CD63 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Number	MEM-259
Form/Format	Purified Purified IgG - liquid
Concentration	IgG concentration 1.0 mg/ml (varies by lot)
Sequence Length	238
Applicable Applications for anti-CD63 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Perservative Stabilisers	0.09% Sodium Azide Preparation
Buffer Solution	Phosphate buffered saline
Target Species	Human
Preparation and Storage	Store at 4 degree C or at -20 degree C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life: 18 months from date of despatch.


IMAGES


Application Data

Staining of human peripheral blood monocytes with Mouse anti Human CD63


Application Data

Staining of human peripheral blood monocytes with Mouse anti Human CD63:FITC


Application Data

Peripheral Human Monocytes stained with Mouse anti Human CD63:APC

ADDITIONAL INFORMATION

Related Product Information for anti-CD63 antibody	Mouse anti Human CD63 antibody, clone MEM-259 recognizes the CD63 cell surface antigen, also known as granulophysin, lysosomal associated membrane protein 3 (LAMP-3), Melanoma-associated antigen ME491, Ocular melanoma-associated antigen or tetraspanin-30. CD63 is a 238 amino acid, tetraspanin glycoprotein of ~40-60kDa. CD63 is expressed by monocytes, macrophages, endothelial cells where it is a component of Weibel-Palade bodies (Vischer et al. 1993) and by a variety of cell lines. It is located intracellularly in lysosomal granules of platelets, being translocated to the surface upon activation. It may therefore be useful as a marker of platelet activation (Hamamoto et al. 1994). CD63 expression is also associated with melanoma development (Radford et al. 1997) and expression is rapidly down-regulated by progesterone in endometrial tissues (Okada et al. 1999).
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	383872447
NCBI GeneID	967
NCBI Accession #	NP_001244318.1
NCBI GenBank Nucleotide #	NM_001257389.1
UniProt Accession #	P08962; Q5TZP3; Q8N6Z9; Q9UCG6; F8VZE2
Molecular Weight	17,271 Da
NCBI Official Full Name	CD63 antigen isoform A
NCBI Official Synonym Full Names	CD63 molecule
NCBI Official Symbol	CD63
NCBI Official Synonym Symbols	MLA1; ME491; LAMP-3; OMA81H; TSPAN30
NCBI Protein Information	CD63 antigen; CD63 antigen (melanoma 1 antigen); granulophysin; lysosomal-associated membrane protein 3; lysosome-associated membrane glycoprotein 3; melanoma-associated antigen ME491; melanoma-associated antigen MLA1; ocular melanoma-associated antigen; tetraspanin-30; tspan-30
UniProt Protein Name	CD63 antigen
UniProt Gene Name	CD63
UniProt Synonym Gene Names	MLA1; TSPAN30; LAMP-3; Tspan-30
UniProt Entry Name	CD63_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd63-antibody-clone-mem-259-immunoprecipitation-immunofluorescence-flow-cytometry-elisa/49336>