

Rat CD49f Monoclonal Antibody | anti-CD49f antibody

RAT ANTI HUMAN CD49f

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA49430	Rat	Cynomolgus monkey, Dog, Mouse, Pig, Sheep	Immunoprecipitation, Flow Cytometry, Functional Assay, Immunohistochemistry


PRODUCT SYNONYMS

CD49f; Monoclonal Antibody; RAT ANTI HUMAN CD49f; anti-CD49f antibody


PRODUCT SPECIFICATIONS

Host	Rat
Reactivity	Cynomolgus monkey, Dog, Mouse, Pig, Sheep
Clonality	Monoclonal
Isotype	IgG2a
Clone Number	NKI-GoH3
Form/Format	Purified Purified IgG - liquid
Concentration	IgG concentration 1.0 mg/ml (varies by lot)
Sequence Length	1073
Applicable Applications for anti-CD49f antibody	IP (Immunoprecipitation), FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry)
Perservative Stabilisers	0.09% Sodium Azide Preparation
Histology Positive Control Tissue	Human Tonsil Buffer Solution
Target Species	Human
Preparation and Storage	Store at 4 degree C or at -20 degree C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life: 18 months from date of despatch.

IMAGES



Application Data
Staining of human peripheral blood platelets with Rat anti Human CD49f:Biotin



Application Data
Staining of human peripheral blood lymphocytes with Rat anti Human CD49f:Biotin

ADDITIONAL INFORMATION

Related Product Information for anti-CD49f antibody	Rat anti Human CD49f antibody, clone NKI-GoH3 recognizes CD49F (VLA-6 alpha chain), a 120kD cell surface glycoprotein that can form distinct complexes with CD29 (VLA beta-chain), resulting in the VLA-6 (alpha-6 beta-1) complex, which is expressed on human platelets, or with the beta-4 integrin resulting in the alpha-6 beta-4 complex which is expressed on various human epithelial cells. Clone NKI-GoH3 reacts with platelets, megakaryocytes, T lymphocytes and common Acute Lymphoblastic Leukaemia cells (alpha-6 beta-1). In immunohistology the monoclonal antibody reacts with epithelial cells of a variety of tissues, peripheral nerves, microvascular endothelial cells, placenta cyto- and syncytiotrophoblasts.VLA-6 is an important mediator of cell binding to laminin. Studies suggest that clone NKI-GoH3 specifically blocks the binding of cells to the E8 fragment of laminin (Sonnenberg et al. 1998).
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	119395742
NCBI GeneID	3655
NCBI Accession #	NP_000201.2
NCBI GenBank Nucleotide #	NM_000210.2
UniProt Accession #	P23229; Q08443; Q0MRC7; Q14646; Q16508; Q53RX7; Q59HB7; B2RMU9; B4DG69; B4DKB8; C4AM96; G5E9H1
Molecular Weight	122,358 Da
NCBI Official Full Name	integrin alpha-6 isoform b
NCBI Official Synonym Full Names	integrin, alpha 6
NCBI Official Symbol	ITGA6
NCBI Official Synonym Symbols	CD49f; VLA-6; ITGA6B
NCBI Protein Information	integrin alpha-6; CD49 antigen-like family member F; integrin alpha6B
UniProt Protein Name	Integrin alpha-6
UniProt Gene Name	ITGA6
UniProt Synonym Gene Names	Alpha6p
UniProt Entry Name	ITA6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd49f-antibody-clone-nki-goh3-immunoprecipitation-flow-cytometry-immunohistochemistry-elisa/49430>